



# جامعة ستاردوم

مجلة ستاردوم العلمية للدراسات الهندسية و الطبيعية




— مجلة ستاردوم العلمية للدراسات الهندسية و الطبيعية —

تصدر بشكل نصف سنوي عن جامعة ستاردوم

المجلد الثاني ا العدد الأول- لعام 2024م

رقم الإيداع الدولي : ISSN 2980-3756





**Peer Reviewing Committee of Stardom Journal  
for Natural and Engineering Sciences**

**Head Of Committee**

Prof. Taha Elwi - Iraq

**Manager Of Committee**

Prof. Abbas Al-Azzawi - Iraq

**Language Editor**

Mrs. Laila Hussein Al-Eyan - Turkey

**Member of Committee**

Prof. Ali Saegh - Iraq

Eng. Saeed Ahmed - Philistine

Prof. Hayder H. Al-Khaylani - Iraq

Dr. Salem Saleh Barahmah - Yemen

Prof. Ali Abdulmajeed Ali Ahmed - Yemen

Prof. Mohammed Moqbil - Yemen

**All literary and artistic property rights are reserved to  
Stardom Journal for Engineering and Natural Sciences**

- ▶ Unlocking the Cholesterol Content: A Comprehensive Analysis of Cholesterol Levels in Dairy Products, Fats, and Oils Through High-Performance Liquid Chromatography

Mokhtar S. S. Al - Salimi-Fares S. S. Al-Saidi - Adel A. M. Saeed

- ▶ Bilateral Generating Functions for the Two-Parameter, Three-variable Srivastava polynomials

Dr. Salem Saleh Barahmah

- ▶ A new extended beta function involving generalized mittag-leffler function and it's Applications

Dr. Salem Saleh Barahmah

▶ تربية و زراعة بعض أنواع النحل البري الملقح لطيف واسع من النباتات

أ.د. عبدالسلام محمد - Prof:Abdoalsalam mohamed gaool Al-Hjry

▶ مقال بحثي في كيمياء تحليل البيئة

دراسة بعض الصفات الفيزيوكيميائية والملوثات غير العضوية للمياه العادمة الناتجة من مدبغة لودر للبيئة المجاورة

جمال أحمد عبدالله الدهبلي - علي ناصر أحمد الكوم - عادل أحمد محمد سعيد

- ▶ Flora Abyan governorate

Abdul-Nasser Al-Gefrli

- ▶ Investigation of the Absorbing the Crash Impact of Car Accedient Due Tube Inversion

Eng. Abdulhakim Hamood Ahmed Abdulwahid-Dr. Fawaz Ahmed Ghaleb

Noman-Eng. Nezar Nasser Ali Haithm

- ▶ Investigation of the inside temperature to arrive comfortable condition1: case study Mechanical workshop building

Dr. Fawaz Ahmed Ghaleb Noman

# **Stardom Journal for Natural and Engineering Sciences (SJNES)**

## **Introduction**

Aims and Scope:

The stardom Journal for Natural and Engineering sciences is a journal for the publication of peer reviewed, original research for all aspects of Natural Sciences and Engineering Sciences.

This journal provides a forum for the discussion of Natural and Engineering sciences problems around the world and for the presentation of results. It is aimed only at the Natural and Engineering sciences.

Research Areas Include, but are not exclusive to:

- Biology and biomedical engineering
- Environmental Sciences
- Chemistry studies and Chemical Engineering
- Physics studies
- Material Studies
- Mathematics
- Computer Sciences and engineering
- Civil Engineering
- Area Engineering
- Environmental Engineering
- Telecommunications Engineering and networking

- Computer Engineering
- Electrical Engineering
- Agriculture Engineering
- Human-machine systems
- Space exploration
- Renewable energy
- Power systems and control
- Artificial intelligence and its applications
- Theoretical studies

Papers submitted should address Stardom Natural and Engineering sciences issues using a range of techniques e.g. case studies, observational and theoretical analyses, the application of science, engineering and technology to questions of environmental concern or mathematical and computer modeling techniques with the aim of informing both the researcher and practitioner.

### **Before you Begin**

#### **Conflict of interest**

All authors are requested to disclose any actual or potential conflict of interest including any financial, personal or other relationships with other people or organizations within three years of beginning the submitted work that could inappropriately influence, or be perceived to influence, their work.

#### **Submission declaration and verification**

Submission of an article implies that the work described has not been published previously (except in the form of an abstract or as part of a published lecture or academic thesis or as an electronic preprint, that it is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyright-holder.

#### **Changes to authorship**

This policy concerns the addition, deletion, or rearrangement of author names in the authorship of accepted manuscripts:

Before the accepted manuscript is published in an online issue: Requests to add or remove an author, or to rearrange the author names, must be sent to the Journal Manager from the corresponding author of the accepted manuscript and must include: (a) the reason the name should be added or removed, or the author names rearranged and (b) written confirmation (e-mail, fax, letter) from all authors that they agree with the addition, removal or rearrangement. In the case of addition or removal of authors, this includes confirmation from the author being added or removed. Requests that are not sent by the corresponding author will be forwarded by the Journal Manager to the corresponding author, who must follow the procedure as described above. Note that: (1) Journal Managers will inform the Journal Editors of any such requests and (2) publication of the accepted manuscript in an online issue is suspended until authorship has been agreed.

After the accepted manuscript is published in an online issue: Any requests to add, delete, or rearrange author names in an article published in an online issue will follow the same policies as noted above and result in a corrigendum..

### **Role of the funding source**

You are requested to identify who provided financial support for the conduct of the research and/or preparation of the article and to briefly describe the role of the sponsor(s), if any, in study design; in the collection, analysis and interpretation of data; in the writing of the report; and in the decision to submit the article for publication. If the funding source(s) had no such involvement then this should be stated.

### **Language**

Please write your text in good English (American or British usage is accepted, but not a mixture of these). Authors who feel their English language manuscript may require editing to eliminate possible grammatical or spelling errors and to conform to correct scientific English may wish to use the English Language Editing, as well as Arabic Language is accepted.

### **Submission**

Submission to this journal proceeds totally online and you will be guided stepwise through the creation and uploading of your files. The system automatically converts source files to a single PDF file of the article, which is used in the peer-review process. Please note that even though manuscript source files are converted to PDF files at submission for the review process, these source files are needed for further processing after acceptance.

## **Referees**

Please submit, with the manuscript, the names and addresses of at least three potential referees.

## **Preparation**

### **Use of word-processing software**

It is important that the file be saved in the native format of the word processor used. The text should be in single-column format. Keep the layout of the text as simple as possible. Most formatting codes will be removed and replaced on processing the article. In particular, do not use the word processor's options to justify text or to hyphenate words. However, do use bold face, italics, subscripts, superscripts etc. When preparing tables, if you are using a table grid, use only one grid for each individual table and not a grid for each row. If no grid is used, use tabs, not spaces, to align columns. The electronic text should be prepared in a way very similar to that of conventional manuscripts.

Note that source files of figures, tables and text graphics will be required whether or not you embed your figures in the text.

To avoid unnecessary errors, you are strongly advised to use the 'spell-check' and 'grammar-check' functions of your word processor.

Please **do not** add line numbering to the text document.

## **Article structure**

### **Subdivision - numbered sections**

Divide your article into clearly defined and numbered sections. Subsections should be numbered 1.1 (then 1.1.1, 1.1.2, ...), 1.2, etc. (the abstract is not included in section numbering). Use this numbering also for internal cross-referencing: do not just refer to 'the text'. Any subsection may be given a brief heading. Each heading should appear on its own separate line.

## **Introduction**

State the objectives of the work and provide an adequate background, avoiding a detailed literature survey or a summary of the results.

## **Material and methods**

Provide sufficient detail to allow the work to be reproduced. Methods already published should be indicated by a reference: only relevant modifications should be described.

## **Results**

Results should be clear and concise.

## **Discussion**

This should explore the significance of the results of the work, not repeat them. A combined Results and Discussion section is often appropriate. Avoid extensive citations and discussion of published literature.

## **Conclusions**

The main conclusions of the study must be presented in a short Conclusions section, which may stand alone or form a subsection of a Discussion or Results and Discussion section.

## **Essential title page information**

- **Title.** Concise and informative. Titles are often used in information-retrieval systems. Avoid abbreviations and formulae where possible.
- **Author names and affiliations.** Where the family name may be ambiguous (e.g., a double name), please indicate this clearly. Present the authors' affiliation addresses (where the actual work was done) below the names. Indicate all affiliations with a lower-case superscript letter immediately after the author's name and in front of the appropriate address. Provide the full postal address of each affiliation, including the country name and, if available, the e-mail address of each author.
- **Corresponding author.** Clearly indicate who will handle correspondence at all stages of refereeing and publication, also post-publication. **Ensure that phone numbers (with country and area code) are provided in addition to the e-mail address and the complete postal address. Contact details must be kept up to date by the corresponding author.**
- **Present/permanent address.** If an author has moved since the work described in the article was done, or was visiting at the time, a 'Present address' (or 'Permanent address') may be indicated as a footnote to that author's name. The address at which the author actually did the work must be retained as the main, affiliation address. Superscript Arabic numerals are used for such footnotes.

## **Abstract**

A concise and factual abstract is required. The abstract should state briefly the purpose of the research, the principal results and major conclusions. An abstract is often presented separately from the article, so it must be able to stand alone. For this reason, References should be avoided, but if essential, then cite the author(s) and year(s). Also, non-standard or uncommon abbreviations should be avoided, but if essential they must be defined at their first mention in the abstract itself. Not more 250 words.

## **Graphical abstract**

A Graphical abstract is optional and should summarize the contents of the article in a concise, pictorial form designed to capture the attention of a wide readership online. Authors must provide images that clearly represent the work described in the article. Graphical abstracts should be submitted as a separate file in the online submission system. Image size: Please provide an image with a minimum of  $531 \times 1328$  pixels (h  $\times$  w) or proportionally more. The image should be readable at a size of  $5 \times 13$  cm using a regular screen resolution of 96 dpi. Preferred file types: TIFF, EPS, PDF or MS Office files.

## **Highlights**

High lights are mandatory for this journal. They consist of a short collection of bullet points that convey the core findings of the article and should be submitted in a separate file in the online submission system. Please use 'Highlights' in the file name and include 3 to 5 bullet points (maximum 85 characters, including spaces, per bullet point).

## **Keywords**

Immediately after the abstract, provide a maximum of 6 keywords, using American spelling and avoiding general and plural terms and multiple concepts (avoid, for example, 'and', 'of'). Be sparing with abbreviations: only abbreviations firmly established in the field may be eligible. These keywords will be used for indexing purposes.

## **Acknowledgements**

Collate acknowledgements in a separate section at the end of the article before the references and do not, therefore, include them on the title page, as a footnote to the title or otherwise. List here those individuals who provided help during the research (e.g., providing language help, writing assistance or proof reading the article, etc.).

## **Math formulae**

Present simple formulae in the line of normal text where possible and use the solidus (/) instead of a horizontal line for small fractional terms, e.g., X/Y. In principle, variables are to be presented in italics. Powers of e are often more conveniently denoted by exp. Number consecutively any equations that have to be displayed separately from the text (if referred to explicitly in the text).

## **Footnotes**

Footnotes should be used sparingly. Number them consecutively throughout the article, using superscript Arabic numbers. Many word processors build footnotes into the text, and this feature may be used. Should this not be the case, indicate the position of footnotes in the text and present the footnotes themselves separately at

the end of the article. Do not include footnotes in the Referencelist.  
Table footnotes

Indicate each footnote in a table with a superscript lowercase letter.

## **Artwork**

### **Electronic artwork**

#### General points

- Make sure you use uniform lettering and sizing of your original artwork.
- Save text in illustrations as 'graphics' or enclose the font.
- Only use the following fonts in your illustrations: Arial, Courier, Times, Symbol.
- Number the illustrations according to their sequence in the text.
- Use a logical naming convention for your artwork files.
- Provide captions to illustrations separately.
- Produce images near to the desired size of the printed version.
- Submit each figure as a separate file.

#### Formats

Regardless of the application used, when your electronic artwork is finalized, please 'save as' or convert the images to one of the following formats (note the resolution requirements for line drawings, halftones, and line/halftone combinations given below):

EPS: Vector drawings. Embed the font or save the text as 'graphics'.

TIFF: Color or grayscale photographs (halftones): always use a minimum of 300 dpi.

TIFF: Bitmapped line drawings: use a minimum of 1000 dpi.

TIFF: Combinations bitmapped line/half-tone (color or grayscale): a minimum of 500 dpi is required.

If your electronic artwork is created in a Microsoft Office application (Word, PowerPoint, Excel) then please supply 'as is'.

#### **Please do not:**

- Supply files that are optimized for screen use (e.g., GIF, BMP, PICT, WPG); the resolution is too low;
- Supply files that are too low in resolution;
- Submit graphics that are disproportionately large for the content.

#### **Color artwork**

Please make sure that artwork files are in an acceptable format (TIFF, EPS or MS Office files) and with the correct resolution. If, together with your accepted. Please note: Because of technical complications which can arise by converting color figures to 'gray scale' (for the printed version should you not opt for color in print) please submit in addition usable black and white versions of all the color illustrations.

### **Figure captions**

Ensure that each illustration has a caption. Supply captions separately, not attached to the figure. A caption should comprise a brief title (**not** on the figure itself) and a description of the illustration. Keep text in the illustrations themselves to a minimum but explain all symbols and abbreviations used.

### **Tables**

Number tables consecutively in accordance with their appearance in the text. Place footnotes to tables below the table body and indicate them with superscript lowercase letters. Avoid vertical rules. Be sparing in the use of tables and ensure that the data presented in tables do not duplicate results described elsewhere in the article.

### **References**

#### **Citation in text**

Please ensure that every reference cited in the text is also present in the reference list (and vice versa). Any references cited in the abstract must be given in full. Unpublished results and personal communications are not recommended in the reference list, but may be mentioned in the text. If these references are included in the reference list they should follow the standard reference style of the journal and should include a substitution of the publication date with either 'Unpublished results' or 'Personal communication'. Citation of a reference as 'in press' implies that the item has been accepted for publication.

#### **Web references**

As a minimum, the full URL should be given and the date when the reference was last accessed. Any further information, if known (DOI, author names, dates, reference to a source publication, etc.), should also be given. Web references can be listed separately (e.g., after the reference list) under a different heading if desired, or can be included in the reference list.

#### **References in a special issue**

Please ensure that the words 'this issue' are added to any references in the list (and any citations in the text) to other articles in the same Special Issue.

## Reference style

**Text:** Indicate references by number(s) in square brackets in line with the text. The actual authors can be referred to, but the reference number(s) must always be given. Example: '..... as demonstrated [3,6]. Barnaby and Jones [8] obtained a different result ....'

**List:** Number the references (numbers in square brackets) in the list in the order in which they appear in the text.

**Examples:**

Reference to a journal publication:

[1] J. van der Geer, J.A.J. Hanraads, R.A. Lupton, The art of writing a scientific article, *J. Sci. Commun.* 163 (2010) 51–59.

Reference to a book:

[2] W. Strunk Jr., E.B. White, *The Elements of Style*, fourth ed., Longman, New York, 2000.

Reference to a chapter in an edited book:

[3] G.R. Mettam, L.B. Adams, How to prepare an electronic version of your article, in: B.S. Jones, R.Z. Smith (Eds.), *Introduction to the Electronic Age*, E-Publishing Inc., New York, 2009, pp. 281–304.

## Submission checklist

The following list will be useful during the final checking of an article prior to sending it to the journal for review. Please consult this Guide for Authors for further details of any item.

### Ensure that the following items are present:

One author has been designated as the corresponding author with contact details:

- E-mail address
- Full postal address
- Phone numbers

All necessary files have been uploaded, and contain:

- Keywords
- All figure captions

- All tables (including title, description, footnotes)

Further considerations

- Manuscript has been 'spell-checked' and 'grammar-checked'
- References are in the correct format for this journal
- All references mentioned in the Reference list are cited in the text, and vice versa
- Permission has been obtained for use of copyrighted material from other sources

# **Unlocking the Cholesterol Content: A Comprehensive Analysis of Cholesterol Levels in Dairy Products, Fats, and Oils Through High-Performance Liquid Chromatography**

**Mokhtar S. S. Al-Salimi<sup>1</sup>**

**Fares S. S. Al-Saidi<sup>1</sup>**

**Adel A. M. Saeed<sup>2\*</sup>**

**<sup>1</sup>Department of Chemistry/Yafea University College, University of Lahej,  
Lahej, Yemen.**

**<sup>2\*</sup>Department of Chemistry/Faculty of Science, University of Aden,  
Khormaksar, Aden, Yemen, Mob. no. &**

## **Abstract**

This study aimed to evaluate the cholesterol content in various dairy products commonly consumed by the general population, as well as fats and oils used in cooking and food preparation. The samples were collected from local markets and farmers in the Aden and Abyan Governorates. To analyze the samples, a highly sensitive high-performance liquid chromatography (HPLC) instrument was utilized. The findings of this study indicated that the highest cholesterol levels were observed in samples of animal ghee and hydrogenated vegetable ghee. Additionally, dairy products derived from full-fat animal milk displayed higher cholesterol content. Specifically, Bint ghee had the highest cholesterol percentage (495.0 mg/100g), followed by local ghee (b) (481.5 mg/100g), Qamaria ghee (453.48 mg/100g), Amazon ghee (411.83 mg/100g), Jabli ghee (368.14 mg/100g), and local ghee (a) (304 mg/100g). Butter samples had a cholesterol concentration of 245 mg/100g, while sheep cheese contained 84.3 mg/100g. Areej ghee recorded 51.9 mg/100g, and bovine cheese had 49.3 mg/100g. The cholesterol percentage in Saudi cheddar cheese reached 42.0 mg/100g. On the other hand, cream, margarine, sesame oil, and Shifa oil samples had cholesterol concentrations below the detection limits, indicating minimal or no cholesterol content.

**Keywords:** Cholesterol, Dairy products, Fats and oils, High-performance liquid chromatography

## **1. Introduction**

Cholesterol, a lipid-like compound, plays a crucial role in various physiological processes within the human body [1,2]. It is present in cell membranes, the brain, nervous tissues, and the myelin sheath. Cholesterol is also found in substantial amounts in the liver, bile salts, and the skin, where exposure to sunlight converts it into vitamin D. Moreover, the adrenal gland utilizes cholesterol for synthesizing steroid hormones like cortisol, estrogen, and progesterone. The liver produces cholesterol by utilizing fats, carbohydrates, and proteins [1]. Additionally, dietary sources, such as meats, milk, and eggs, contribute to excess cholesterol in the body [3-7]. Conversely, vegetables and plant-based products do not contain cholesterol. The average American daily diet typically contains 400-500 mg of cholesterol, the highest intake worldwide, although the American Heart Association recommends intake of no more than 300 mg per day [3]. Research suggests a link between saturated fats, cholesterol, and various diseases including diabetes, breast cancer, pancreatic cancer, colon cancer, and atherosclerosis. Atherosclerosis occurs due to

the buildup of fatty protein compounds in blood vessels, leading to restricted blood flow, thereby increasing the risk of heart attacks. Factors like lack of exercise, smoking, obesity, and aging further compound the risk of heart disease. Clinically, high cholesterol levels are defined as total cholesterol levels in plasma exceeding 200 mg [8-10]. In the diet, saturated fats can trigger cholesterol production in the liver. Thus, adopting a diet low in cholesterol and saturated fats can help lower blood cholesterol levels. The American Institute for Cancer Research recommends increasing fiber and starch intake through the consumption of vegetables, fruits, and grains, along with moderate portions of low-fat, low-cholesterol foods such as fish, poultry, and lean meats. It is also advised to limit the consumption of cholesterol-rich foods, including eggs, nuts, fried potatoes, red meats, and full-fat dairy products [1]. This study aimed to estimate the cholesterol content in specific dairy products, fats, and local and imported oils. The objective was to verify the cholesterol levels in these products, which are considered significant dietary sources of cholesterol and fats, and to compare them with studies conducted in several countries, utilizing High-Performance Liquid Chromatography (HPLC) as the analytical method.

## **2. Materials and Research Methods**

A Japanese-made high-performance liquid chromatography (HPLC) instrument, specifically the JASCO Co-2065 Plus model, was utilized in this experiment. The instrument comprised a Quaternary Gradient Pump (JASCO PU-2089 Plus) and an Intelligent UV/Vis Detector (JASCO UV-2070 Plus) capable of operating in the visible and ultraviolet ray range (UV-VIS). These instrument components were connected to a computer that housed the necessary programs for the analysis.

For the separation process, a C8 column (Mediterranea HPLC, Mediterranea C8 5 $\mu$ m, 15cm x 0.46cm) was employed, which contained Octyl Silyl (OS) (-Si-CH<sub>2</sub>)<sub>7</sub>-CH<sub>3</sub>) packing material. This column was accommodated in an Intelligent Column Oven (JASCO LC-Net II/ADC).

### **2-1 Collection of samples**

Samples were obtained from various sources including the local market and farmers involved in livestock breeding in the Aden and Abyan governorates. The collection of samples was carried out in a random manner to ensure representative sampling. These samples were carefully placed in sterilized plastic bags and glass containers to maintain their integrity. Afterward, they were transported to the laboratory for further analysis.

In the laboratory, each individual sample was prepared by undergoing a mixing process, and in some cases, heating was applied to facilitate the homogenization of their components. This step was crucial in order to obtain a representative sample that was blended and homogeneous. Sufficient volume of each sample was taken into consideration to ensure accurate analysis, while also allowing for the possibility of repeating certain analyses.

Following the preparation process, the samples were transferred to clean glass containers and stored in a refrigerator to maintain their stability and prevent any degradation prior to analysis.

### **2-2 Types of samples under study**

**Assorted Dairy Product Samples:** Saudi Cheddar cheese, Five Star cream .1 packaged in the UAE, Yemeni vegetable margarine obtained from Dhamran supermarket in Mansoura-Aden, traditional cow's milk cheese, traditional sheep's milk cheese obtained from Sheikh Othman market in Aden, light yogurt from Al-Hana, Yemeni-made yogurt from Al-Hana, fresh light yogurt traditionally made from cow's milk, and cow's milk butter obtained from Al-Hisn area in Abyan .Governorate

**Ghee and Oil Samples:** Areej ghee made in Oman, Amazon ghee packaged in .2 ,Malaysia, Qamaria ghee, Bint ghee, Jabali ghee, all locally made in Yemen obtained from Dhamran supermarket in Aden governorate. Locally made ghee (a) and (b) made from traditional cow's milk butter at home, sesame oil pressed at the Wali press in Sheikh Othman-Aden, cooking oil (Shifa) made in Oman, obtained .from Al-Hisn area in Abyan

### **2-3 Preparation of the original cholesterol solution**

The primary cholesterol solution was initially prepared by dissolving 0.1 g of pure cholesterol in an adequate amount of methanol in a 100 ml volumetric flask. The solution was then further diluted with methanol to reach the marked level, resulting in an initial stock solution of 1,000 ppm [11].

#### **2-4 Preparation standard solutions from the original cholesterol solution**


To create standard solutions with varying concentrations (10, 20, 30, 40, 50 ppm), the original cholesterol standard solution (1000 ppm) was used. These standard solutions served as reference samples, with 20  $\mu$ L of each solution individually injected into the HPLC.

### **2-5 Calibration curve**

To generate a linear calibration curve, standard solutions were prepared and injected into an HPLC/UV instrument using a mobile phase consisting of acetonitrile,

methanol, and 2-propanol in a ratio of 1:3:7. The injections were done sequentially at a flow rate of 1.6 ml/min and at room temperature. A C8 column was used, and the absorbance was measured at a wavelength of 205 nm. The concentrations of the standard solutions were plotted on the x-axis, while the corresponding response values (peak area) were plotted on the y-axis. The slope and intercept of the calibration curve were determined based on the calculated values, which can be found in Table 1 and Figure 1. The response measurements were repeated three times (n=3), and a strong linear correlation was observed with a correlation coefficient (R<sup>2</sup>) of 0.9986. The linear equation derived from the data is  $y = 5886.33x + 2771.9$  [11].

**Table 1: Linear functions of the standard curve for cholesterol.**



\*It is the lowest detectable concentration of the analyte that can be measured with acceptable statistical accuracy.

<sup>φ</sup>The minimum detectable concentration for the analyte can be estimated using appropriate standard materials.

### Figure 1: Cholesterol calibration curve.

Table 2: Average peak areas for different concentrations of standard solutions of cholesterol.

N O.	Concentration (ppm)	Peak area (micro volts/s) n=3	Wavelength ( $\lambda$ )	Flow rate (mL/min)	Retention time (min)	Mobil Phase
1	10	64680	205 nm	1.6	3.8	(Acetonitrile, Methanol, 2-Propanol) (1:3:7)
2	20	118486				
3	30	178375				
4	40	234057				
5	50	301211				

#### 2-6 Preparation of Saponification Solution

To create the saponification solution, a precise amount of 5.6 g of potassium hydroxide (KOH) was weighed and transferred to a standard 200 ml flask. Methanol was then added to the flask until it reached the mark, resulting in an alcoholic potassium hydroxide solution with a concentration of 0.5 M, as described in references [12,13].

#### 2-7 Preparation of the Moving Phase

The mobile phase utilized in this study consisted of a mixture of acetonitrile, methanol, and 2-propanol in a ratio of 7:3:1. To prepare the mobile phase, 350 ml of acetonitrile, 150 ml of methanol, and 50 ml of 2-propanol were measured using a calibrated standard flask and combined in a 1000 ml bottle. Prior to using the mobile phase in the HPLC instrument, the bottle containing the mixture was subjected to 15 minutes of ultrasonic cleaning to eliminate any trapped gases [14].

#### 2-8 Estimation of Cholesterol in the Studied Samples

1. To extract cholesterol from the dairy product samples, the direct saponification method was employed. Accurate weights of 0.2 g of butter, 0.6 g of cheeses, and 1 g of cream and yogurt were separately transferred to spiral-capped Teflon-coated test tubes. Saponification was carried out by adding 2 ml of 0.5 M methanolic potassium hydroxide solution to each tube, which was then placed in a hot water bath at 80°C. The tube contents were stirred every 5 minutes during a 15-minute saponification period. The unsaponified portion was then extracted by adding 5 ml of hexane and 2 ml of distilled water. After vigorous shaking for 10 minutes, the

mixture was centrifuged at 1400 revolutions per minute for 5 minutes. This extraction process with hexane was repeated three times using 5 ml each time. The resulting hexane extracts were combined in a 50 ml round-bottomed glass flask with a cover. The total extract was evaporated to dryness at 70°C using a rotary evaporator. The remaining extract was dissolved in 2 ml of methanol, and 20 µL of the solution was directly injected into the HPLC instrument. This injection process was repeated three times, and cholesterol quantification was performed using a calibration curve of standard cholesterol solutions. These analyses were conducted at the Higher Institute for Drug Standards and Quality in Khormaksar, Aden Governorate [12,13].

2. The analysis of ghee and oil samples was performed using two different methods. In the first method, local ghee, fragrant ghee, sesame oil, and palm oil underwent direct saponification followed by cholesterol extraction from the unsaponifiable portion using hexane solvent. This extraction process was conducted following the same steps mentioned earlier for the dairy samples. Subsequently, 20 µL of the extract was taken after drying and dissolving it in 2 ml of methanol, which was then injected into an HPLC device. The injection process was repeated three times for consistency [11].

In the second method, samples of local ghee, Qamaria ghee, pent ghee, mountain ghee, and Amazon ghee were prepared by weighing 0.250 g of ghee and dissolving it in 5 ml of acetone. The solution was then transferred to a 50 ml volumetric flask and completed to volume with methanol. To ensure complete dissolution and removal of gases, the flask was subjected to ultrasonic waves for 10 minutes. A portion of the solution was then centrifuged at 1500 revolutions per minute for 10 minutes. Subsequently, 20 µL of the solution was injected into the HPLC instrument three times to obtain consistent readings [11].

### **2-9 Statistical analysis**

The Origin 7.5 program, a powerful statistical analysis tool, was employed to perform rigorous data analysis in this study. To investigate the variations in the data, one-way ANOVA analysis was conducted, which is a robust statistical method commonly utilized in research to determine whether there are significant differences between groups. In this study, a significance level of  $P < 0.05$  was adopted, implying that any observed differences between the means had to be statistically significant at this threshold to be considered meaningful.

To further ascertain the presence of statistically significant differences between the means, the least significant difference (LSD) was calculated. The LSD method is an effective post hoc test that enables researchers to compare pairs of means within the groups and determine if there are any statistically significant disparities. This approach provides a comprehensive understanding of the data by assessing the magnitude and significance of differences between multiple sets of mean values.

By employing the Origin 7.5 program for statistical analysis and conducting one-way ANOVA with a significance level of  $P < 0.05$ , coupled with the calculation of the LSD, this study ensures a rigorous examination of the data, allowing for precise conclusions to be drawn. The implementation of these statistical techniques contributes to the credibility and reliability of the research findings, enabling a comprehensive evaluation of the results obtained [15].

### **3- Results and Discussion**

The chromatographic analysis results of the samples under study are presented in Figures 2-15. These results aim to differentiate between the various types of samples. In particular, the first variety corresponds to dairy products samples, which are classified based on their manufacturing origin and the type of fat added to the product. This includes products with animal fat such as local production butter, cow cheese, sheep cheese, as well as foreign production cheddar cheese. Additionally, there are products with vegetable fat, such as foreign production cream and national production margarine. Furthermore, there are national production products with animal fat, like local light yogurt, and others with vegetable fat, such as Hana light yogurt and Hana yogurt (Table 3).

Moving on to the second variety (Table 4), it pertains to ghee and oil samples. These samples are categorized according to their country of origin and the type of ghee. The classification is as follows: locally produced cow ghee with animal fat, namely local ghee A and local ghee B. Moreover, there is locally produced refined vegetable ghee, including Qamaria, Bint, and Jabali. Additionally, there is commercially produced refined vegetable ghee, including Arij ghee and Amazon ghee. Lastly there are vegetable oils, such as locally produced sesame oil and foreign-origin cooking oil.

To remove fats and release cholesterol from the samples, the direct saponification process was employed as an initial step. This process has proven to be successful in

recovering and extracting cholesterol using appropriate solvents, thus preventing .any loss of cholesterol [16]

Chromatogram




Figure 2: Chromatogram of butter sample (cow)

Chromatogram




Figure 3: Chromatogram of sheep cheese sample

Chromatogram




Figure 4: Chromatogram of bovine cheese sample

Chromatogram




Figure 5: Chromatogram of cheddar cheese sample

Chromatogram




Figure 6: Chromatogram of natural light yogurt sample

Chromatogram




Figure 7: Chromatogram of commercial Al-Hana light yogurt sample

Chromatogram




Figure 8: Chromatogram of Al-Hana yogurt sample

Chromatogram




Figure 9: Chromatogram of local ghee (a) sample

Chromatogram




Figure 10: Chromatogram of local ghee (b) sample

Chromatogram



Figure 11: Chromatogram of Bint ghee sample




Figure 12: Chromatogram of Qamaria ghee sample




Figure 13: Chromatogram of Jabali ghee sample




Figure 14: Chromatogram of Amazon ghee sample




Figure 15: Chromatogram of Arij ghee sample

### 3-1 Dairy products samples

Based on the results presented in Table 3 and Figure 16, it is evident that there is an inverse relationship between the cholesterol content and the type of fat (animal or vegetable) present in the dairy products under investigation. The samples belonging to the first category, which includes butter and cheeses (cow, sheep, and cheddar) with high animal fat content, as well as one domestic production sample with animal fat (local breed), exhibited the highest levels of cholesterol content (31.45 mg/100g - 245 of the product). Conversely, the commercial samples with removed animal fat and added vegetable fat (Hana light yogurt and Hana yogurt) showcased the lowest cholesterol content, aligning with the aforementioned inverse relationship. Cream and margarine samples had cholesterol levels below the detection limit of the device.

In our study, we observed that the highest concentration of cholesterol in the butter sample (245 mg/100g) exceeded values reported in previous studies [13,17-19] which ranged from 228.1 mg/100g to 235.6 mg/100g. However, our results were in line with the cholesterol content recorded in [20, 21] (240 mg/100g, 244 mg/100g)

Moreover, our findings were lower than values reported as 270.0 mg/100g, 286.4 mg/100g, and 307.0 mg/100g in other studies [22-24]. We also noticed that the cholesterol content in our butter sample closely matched the findings in [14], where the cholesterol content in butter was determined to be 253.5 mg/100g using the same method (HPLC). The study also indicated a cholesterol content of 84.6 mg/100g in .cheese, which aligns with our observation of sheep milk cheese (84.243 mg/100g) However, the cholesterol content in our study for cow's milk cheese and cheddar cheese samples differed. The cholesterol content in the sheep milk cheese sample corresponded to the findings of a study [25] (78.8-97.9 mg/100g). In contrast, the cholesterol content in the butter sample ranged from 227.3 mg/100g to 307.0 ,mg/100g, which matched our study's measurement (245 mg/100g). Furthermore our cheese samples' cholesterol content agreed with [20] (17-114 mg/100g) and was close to [26] (52.3-76.6 mg/100g), while slightly different from [13] (61.9–143.3 mg/100g). Notably, the cholesterol content in our sheep milk cheese (84.24 .mg/100g) aligned with previous studies [13,20,26]

The observed cholesterol levels in our cow's milk cheese (49.13 mg/100g) and cheddar cheese (41.96 mg/100g) were consistent with [20], albeit slightly lower than In study 27, which aimed to determine the cholesterol content in various .[13,26] types of cheese, the values noted for blue cheese (50.38 mg/100g), natural cheese and hard cheese (58.36 mg/100g) were comparable to the findings ,(mg/100g 50.38) in cow's milk cheese and cheddar cheese. However, they were lower than the .cholesterol content in sheep's milk cheese (84.243 mg/100g) in our recent study Additionally, the cholesterol content in goat milk cheese and sheep milk cheese in were 58.61 mg/100g and 65.06 mg/100g, respectively, which were lower than [27] our current study's results. In our study, the cholesterol content in cheddar cheese was lower than the values reported for English cheddar cheese in (mg/100g 41.96) and in both studies [17,19] (107 mg/100g and 102.41 (mg/100g 101.2) [13] .(mg/100g, respectively

Comparing the cholesterol content (mg/100g) in the samples of natural light yogurt Al-Hana injections (4.52 mg/100g), and commercial Al-Hana ,(mg/100g 6.29) yogurt (3.8 mg/100g) with [27] (8.17 mg/100g), we observe that our values are somewhat similar. The cholesterol content recorded in our study for natural light yogurt with animal fat was higher than that in the injections and Al-Hana yogurt with vegetable fat and no animal fat. This explains the lower cholesterol content in .these samples, which is consistent with the cholesterol content in skimmed dairy

Furthermore, [14,20] reported higher cholesterol content in full-fat animal yogurt compared to our (mg/100g, 33.6 mg/100g, and 12.2 mg/100g, respectively 12.46) study's findings, which is expected for light yogurt and Al-Hana yogurt made from animal fat. However, study 15, which employed the HPLC method to determine the cholesterol content in yogurt, reported a cholesterol level of 12.7 mg/100g, slightly higher than our recorded value. Moreover, the cholesterol content reported for yogurt samples (7.33–7.87 mg/100g) in [28] closely aligns with our findings. The cholesterol content in the sample of natural light yogurt was 6.29 mg/100g, while ,yogurt and Al-Hana light yogurt recorded levels of 4.53 mg/100g and 3.8 mg/100g .respectively

The variation in cholesterol content between previous studies and our study may be attributed to various factors, including differences in milk fat content, sample variations, extraction methods, and the cholesterol determination or estimation methods employed.

**Table 3: Cholesterol concentration in selected dairy product samples in mg/100g**

Origin of samples	Type of fat added to the product	Sample Name	Peak area mean	Average concentration (mg/100g)	Standard Deviation (SD)	Relative Standard Deviation (RSD)
Local samples made in a traditional manner.	product animal fat	Butter	1444494	245a	3.357	1.37
		Sheep cheese	1490516	Eighty-four point two four billion	1.226	1.455
		Bovine cheese	870409	49.12c	1.189	2.42
		Cheddar cheese	743768	42nd	1.497	3.57
LSD= 3.078.						
External samples	Animal fat	Local light yogurt	187921	6.3e,g	0.046	0.734
		Hana light yogurt	136062	4.5f,e,g	0.091	2.03
		Hana yogurt	115044	3.81g	0.140	3.66
	LSD = 0.159					
Vegetable fat	Cream (Five Star) Margarine (Momtaz)		N.D			
			N.D			

### 3-2 Samples of ghee and oil

The analysis of clarified butter and oil samples, as presented in Table 4 and Figure 17, yielded interesting findings. The sample of clarified butter labeled "Bint" displayed the highest cholesterol content (495.04 mg/100g) after undergoing dissolution in methanol without direct saponification and subsequent extraction with hexane. Conversely, the vegetable ghee "Areej" exhibited the lowest cholesterol level (51.96 mg/100g).

Regarding sesame oil and palm oil samples, which underwent the same treatment as Areej and local clarified butter (ghee b) (saponification followed by extraction of the unsaponified portion with hexane), their cholesterol content fell below the detection limit of the device (N.D). This observation aligns with existing scientific literature [29], which identifies cholesterol as an animal sterol.

The remaining samples, including local clarified butters (ghee a) made from cow's milk fat, Qamiria fats, Bint, Jabali (locally made), and Amazon (imported), are primarily composed of vegetable oil, mostly palm oil. These products are commonly referred to as vegetable ghee or hydrogenated ghee. The labels attached to them state that they are "cholesterol-free".

To analyze these five samples, a distinct approach was employed, involving the dissolution of 0.250g of each sample in an appropriate amount of acetone. The volume was then completed with methanol up to the mark in a 50 ml volumetric flask, as described in [11]. Subsequent analysis using HPLC-UV revealed the following cholesterol content in these samples: 304 mg/100g, 453.5 mg/100g, 495.04 mg/100g, 368.1 mg/100g, and 411.8 mg/100g respectively.

Comparing the cholesterol content in local clarified butter (a) (304 mg/100g) and local clarified butter (b) (481.5 mg/100g), we note that sample (a) has a higher cholesterol content despite both being local clarified butter. This disparity can be attributed to the different treatment methods employed. Saponification, a process that releases cholesterol from fatty acids bound to it in esters, as well as from cholesterol bound to lipoproteins, significantly contributes to the overall cholesterol content. The subsequent extraction with hexane, an effective solvent for cholesterol extraction, further aids in capturing the released cholesterol. On the other hand, local clarified butter sample (a) only contains free cholesterol since it did not undergo saponification. Consequently, the cholesterol content is limited to free cholesterol, resulting in a lower level compared to sample (b).

This interpretation is corroborated by the striking similarity in cholesterol content between basil ghee (306.5 mg/100g), Pharma ghee (304.1 mg/100g), and Almarai ghee (307.6 mg/100g) as reported in [11]. Moreover, the cholesterol content in sample (b) (304 mg/100g) treated in the same manner is lower than that in sample (a), which underwent saponification and hexane extraction, indicating the effectiveness of these processes in enhancing cholesterol extraction.

The cholesterol content in local clarified butter (a) falls within the range reported in [30] (303 mg/100g to 328 mg/100g), slightly surpasses that in [31] (301 mg/100g), and significantly exceeds the value reported in [32] (256 mg/100g).

Interestingly, the cholesterol content in samples of hydrogenated vegetable ghee (Qamaria, Bint, Jabali, and Amazon) exceeds that in local clarified butter (ghee a), despite undergoing the same treatment. This disparity contradicts the "cholesterol-free" claim on the commercial packaging labels and the assertions made in promotional TV advertisements for these products, as they are supposed to be

cholesterol-free due to their plant origin. Furthermore, the cholesterol content in these studied hydrogenated vegetable ghee samples surpasses the highest observed levels in Mazola ghee (302.8 mg/100g) and Halwani ghee (226.6 mg/100g) [11]. The variation in cholesterol content observed in previous studies compared to our current study may be attributed to sample discrepancies or lack of standardization, ineffective extraction methods employed, differences in fat content in milk, lack of specificity in the estimation methods utilized, and seasonal variations. Additionally, the density of the fat emulsion, and consequently the cholesterol content, may differ, with high-density fat emulsion showing greater cholesterol amounts compared to low-density fat emulsion.

**Table 4: Concentration of cholesterol in samples of ghee and oil in mg/100g**

Sample origin	Type of ghee	Sample Name	Peak area mean	Cholesterol concentration (mg/100g)	Standard deviation (SD)	Relative standard deviation (RSD)
Local	Cow ghee	Local ghee a	92630	304	11.464	3.77
		Local ghee b	569543	481.5	21.505	4.46
Local	Vegetable ghee	Qamaria ghee	136243	453.50	11.11	2.45
	Commercially hydrogenated	Bint ghee	148475	495.0	15.74	3.17
		Jabali ghee	111223	368.14	4.277	1.159
Imported	Vegetable ghee	Arij ghee	63957	51.96	2.365	4.55
	Commercially hydrogenated	Amazon ghee	123984	411.83	1.632	0.395
Local	Vegetable oil	Sesame oil	N.D			
Imported	Vegetable oil	Cooking oil (Shifa)	N.D			
LSD = 14.843						




Figure 16: Cholesterol content graphically in samples of dairy products (mg/100g)




Figure 17: Cholesterol content graphically in ghee samples (mg/100g)




Figure 18: Comparison of cholesterol content in samples of dairy products and ghee (mg/100g)

### 3-3 Comparison of results of samples of dairy derivatives, fats, and oils

In the comparison of cholesterol content in samples of dairy products, ghee, and oil (as presented in Table 5), it is observed that ghee samples exhibit the highest average

cholesterol content at 285.10 mg/100g. On the other hand, dairy product samples show the lowest cholesterol content at 48.32 mg/100g. Previous studies attribute this difference to the animal fat content (suet) present in most animal ghee samples and dairy products, which is higher compared to pure vegetable ghee and dairy products. It is worth noting that there is a direct relationship between the percentage of animal fat and the percentage of cholesterol. In response to the association between consumption of full-fat dairy products and cardiovascular diseases and arteriosclerosis, various food industries have adopted measures recommended by the World Health Organization (WHO). These measures include using dairy products made from skimmed animal milk, low-fat animal milk, and sometimes incorporating vegetable fat into the products.

These adjustments aim to reduce the cholesterol content in the final product, aligning with the WHO's recommendations and addressing potential health concerns associated with the consumption of high-fat dairy products.

**Table 5: Comparison of the average range of cholesterol content in samples of dairy products and ghee mg/100g**

Category	Samples (No.)	Cholesterol concentration (mg/100g)	
		Mean	Range
Dairy Products	9	48.32	N.D - 245
Ghee and Oil	9	285.10	N.D - 495.0

Table 6 and Figure 18 present the cholesterol content in all the studied samples, including dairy products and ghee. The samples of clarified butter exhibited the highest concentration of cholesterol, with values of 495.0 mg/100g, followed by local ghee (b) at 481.5 mg/100g, Qamaria ghee at 453.48 mg/100g, Amazon ghee at 411.83 mg/100g, Jabli ghee at 368.14 mg/100g, and local ghee (a) at 304 mg/100g.

Among the dairy products made from whole animal full-fat milk, the sample of butter displayed a cholesterol content of 245 mg/100g, while sheep milk cheese had a content of 84.3 mg/100g. Arij ghee had a cholesterol content of 51.96 mg/100g, while cow milk cheese had 49.3 mg/100g. Saudi cheddar cheese followed with a value of 42.0 mg/100g.

Samples of cream, margarine, cooking oil (Shifa), and sesame oil were below the detection limits for cholesterol content. Additionally, the samples of raw natural

light yogurt contained 6.2 mg/100g, while the commercial Hana light yogurt had 4.6 mg/100g, and the commercial Hana yogurt had 3.8 mg/100g of cholesterol.


**Table 6: Comparison of cholesterol content in samples of dairy products and ghee in ppm and mg/100g units**

Quality	Type of Fat	Sample Name	Cholesterol Concentration (ppm)	Cholesterol Concentration (mg/100g)
Dairy Products	Whole Animal Full-Fat	Cow butter	2450	245.0
		Sheep cheese	843.3	84.33
		Bovine cheese	490	49.0
		Cheddar cheese	420	42.0
		LSD	5.825	3.078
	Whole Animal Full-Fat	Natural yogurt	62.9	6.29
	Defatted with Added Vegetable Fat	Hana light yogurt	46	4.6
		Hanayogurt	38	3.8
		LSD	0.789	0.159
	Defatted with Added Vegetable Fat	Cream	N.D	N.D
		Margarine	N.D	N.D
	Whole Animal Full-Fat	Local ghee (a)	3040	304
		Local ghee (b)	4815.0	481.50
		LSD	5.356	31.634
Ghee and Oil	National Industry Processed Vegetable Ghee	Bint ghee	4950	495.0
		Qamariaghee	4535.0	453.50
		Jabali ghee	4118.3	411.83
		LSD	0.420	18.072
	Foreign Industry Processed Vegetable Ghee	Amazon ghee	3681.4	368.14
		Arij ghee	519.6	51.96
		LSD	0.591	3.531
	Local Oil	Sesame oil	N.D	N.D
Foreign Oil	Shifa Cooking Oil	N.D	N.D	

### 3-4 Comparison between studied local and imported vegetable ghee samples

Upon comparing local and imported samples, it was observed that local vegetable ghee exhibited higher levels of cholesterol in comparison to imported vegetable ghee. The cholesterol content in local vegetable ghee ranged from 368.14 to 495.0 mg/100g, whereas the cholesterol content in imported vegetable ghee samples varied from 51.96 mg/100g to 411.83 mg/100g. These findings suggest that locally

produced vegetable ghee might have been adulterated with animal fat, as indicated by the results presented in Figure 19 and Table 6. Surprisingly, both local and imported vegetable ghee samples contained cholesterol, despite the product label claiming to be cholesterol-free and derived from hydrogenated palm oil. This discrepancy can be attributed to the lack of regulatory oversight and monitoring by relevant authorities, both for local and imported food products.



**Figure 19:** Cholesterol content in samples of vegetable ghee, animal ghee, local and external oils in mg/100g graphically.

#### 4- Conclusion

In conclusion, the findings of this study highlight significant disparities in the cholesterol content among the analyzed samples. There was a noticeable elevation in cholesterol concentration in samples derived from animal sources in comparison to those obtained from plant sources. Moreover, the results displayed variations in cholesterol content when compared to similar studies conducted in different countries, utilizing standardized analytical techniques such as high-performance liquid chromatography. These dissimilarities may be attributed to various factors, including divergent origins of dairy and fat derivatives in both animal and plant forms, discrepancies in cattle breeds or plant species used for the extraction of ghee and oil, disparities in dietary practices, processing and extraction methods, and the influence of physicochemical and environmental factors. Overall, these findings emphasize the complex nature of cholesterol content in food products and the need for further investigation to comprehend the underlying mechanisms influencing these variations.

## References

- (1) Timberlake Karen. (2016). *An Introduction to General, Organic, and Biological Chemistry*. Twelfth edition. Pearson.533. ISBN-13:978-0-321-90844-5. ISBN-10: 0- 321-90844 -91. Chemistry-Textbooks.
- (2) Li, L., Dutkiewicz, E. P., Huang. Y., Zhou, H., and Hsu, C. (2019). Analytical Methods for Cholesterol Quantification. *Journal of food and drug analysis*, 27, 375-386.
- (3) Campbell Mary, Farrell Shawn, McDougal Owen. (2016). *Biochemistry, Ninth Edition McDougal Student Edition*, 210. ISBN: 978-1-305-96113-5, Loose-leaf Edition:  
ISBN: 978-1-305-96113-1.: 210.
- (4) Gylling, H. (2004). Cholesterol Metabolism and Its Implications for Therapeutic Interventions in Patients with Hypercholesterolaemia. *Int. J. Clin. Pract*, 58, 859–866.
- (5) Derina, K.V., Korotkova, E. I., Dorozhko, E. V., Voronova, O. A. (2017). Voltametric Determination of Cholesterol in Human Blood Serum. *J. Anal. Chem*, 72, 904–910.
- (6) Motonaka, J., Faulkner, L. R. (1993). Determination of Cholesterol and Cholesterol Ester with Novel Enzyme Microsensors. *Anal. Chem*, 65, 3258–3261.
- (7) Saeed, A. A., Salem, T. A., and Al-Saidi, F. S. (2020). Quantitative Determination of the Total Cholesterol in Some Milks of Yemeni Markets. *Electronic Journal of University of Aden for Basic and Applied Sciences*, 12, 111-119.  
<https://doi.org/10.47372/ejua-ba.2020.2.22>.
- (8) Bertolín, J. R., Joy, M., Rufino-Moya, P. J., Lobón, S., Blanco, M. (2018) Simultaneous Determination of Carotenoids, Tocopherols, Retinol and Cholesterol in Ovine Lyophilized Samples of Milk, Meat, and Liver and in Unprocessed/Raw Samples of Fat, *Food Chem*, 257, 182–188.
- (9) Shingla, K. M.; Mehta, B. M. (2018). Cholesterol and Its Oxidation Products: Occurrence and Analysis in Milk and Milk Products. *Int. J. Health Anim. Sci. Food Saf*, 5, 13–39.
- (10) Gorassini, A., Verardo, G.; Fregolent, S. C., Bortolomeazzi, R. Rapid (2017). Determination of Cholesterol Oxidation Products In Milk Powder Based Products by Reversed Phase SPE and HPLC-APCI-MS/MS. *Food Chem*, 230, 604–610.
- (11) AL-Shalan, N. M. (2004). Determination of Cholesterol in Foods, Fats and Blood Plasma by High-Performance Liquid Chromatography. Master Thesis. King Saud University, College Science, Chemistry Department, 7.

- (12) Mohammed, G. A. (2010). Estimation of Cholesterol in Milk by the Enzymatic Method Used to Determine Blood Cholesterol. *Iraq Journal of Agricultural Sciences*, 41(3), 100-105.
- (13) Fletouris, D. J., Botsoglou, N.A., Psomas, I. E., Mantis, A. I.(1998). Rapid Determination of Cholesterol in Milk Products by Direct Saponification and Capillary Gas Chromatography. *J. Dairy Sci*, 81, 2833-2840.
- (14) Oh, H. I., Shin, T. S., Chang, E. J. (2001). Determination of Cholesterol in Milk and Dairy Products by High Performance Liquid Chromatography. *Asian Aust. J. Anim. Sci*, 14(10), 1465-1469.
- (15) AL-Mohammdy, C. (2005). Design and Analysis of Agricultural and Veterinary Experiments. Department of Animal Production and Nutritional Science. Ibb University. Yemen 2005.
- (16) Ulberth, F., and Reich, H. (1992). Gas Chromatographic Determination of Cholesterol in Processed Foods. *Food Chemistry*, 43, 387–391.
- (17) LaCroix, D. E., Mattingly, W. A., Wong, N. P., and Alford, J. A. (1973). Cholesterol, Fat and Protein in Dairy Products. *J. Am. Diet. Assoc*, 62, 275.
- (18) Archard, C. H., Levy, J., and Georgiakakis, N. (1934). Cholesterol des Aliments., *Arch. Mal. Appar. Dig. Mai. Nutr*, 24, 785.
- (19) Man De, J. M., (1964). The Free and Ester Cholesterol Content of Milk and Dairy Products, *Z. Ernaeh*, 5, 1.
- (20) Sieber, R. (1993) Cholesterol Removal from Animal Food- Can it Be justified. Federal Dairy Research institute, CH-3097 Liebefeld (Switzerland).1993.
- (21) Aristova, V. P., and Bekhova, E. K. (1976). Cholesterol in Milk and Milk Products. *Trudy, Vsesoyuznyi Nanchnoissledova-telskii Institut Malochoi Promyshlen no sti*. 42:45 (cf. DSA 1977), 39, 2748.
- (22) Lampert, L. M. (1930). Cholesterol as a Measure of Egg Yolk in Milk Products. *Ind. Eng. Chem. Anal. Ed*, 2, 159.
- (23) Mulder, H., and Zuidhof, T. A. (1958). The Surface Layers of milk Fat Globul. IV. The Cholesterol Content (Free and Esterified Cholesterol) of the Surface Layers. *Netherland Milk Dairy Journal*, 12, 173-179.
- (24) Stewart, D. W. (1923). Some Observations with Regard to Unsaponifiable Matter and Sterols of Edible Fats. *Analyst*, 48,155.
- (25) Sweeny, J. P., and Weihrauch, J. L. (1976). Summary of Available Data for Cholesterol in Food and Methods for its Determination. *Crit. Rev. Food Sci. Nutr*, 8, 131-159.
- (26) Fuke, Y. and Matsuoka, H. (1994). Cholesterol Content and Identification of Foreign Fats in Processed Cheese. *Journal of Japanese Society of Food Nutrition*, 27,269.

- (27) Borkovcova, I., Janouskova, E., Janstonva, M., Drackova, B., and Volrova, L. (2009). Determination of Sterols in Dairy Products and Vegetable Fats by HPLC and GC Methods. *Czech J. Food Sci*, 27, 217-219.
- (28) Tsui, I. C. (1989). Rapid Determination of Total Cholesterol in Homogenized Milk. *J. Assoc. of. Anal. Chem*, 72, 421-424.
- (29) Satyanarayana, U., Chakrapain, U. (2011). *Biochemistry. Book and Allied (P) Ltd. Kolkata, India, P28, 38, 309.*
- (30) Prasad, R., Pandita, N. N. (1990). Cholesterol Content of Milk and Fractionation During Processing. *Indian J. Dairy Sci*, 43, 190–193.
- (31) Kantibhai, M. T. (2007). Development of a method for Cholesterol Estimation in Milk Fat Using Non-Enzymatic Diagnostic Kit". *Master of Science in Dairying, Karnal-132001 (Haryana). India. 2007.*
- (32) Al-Khalifah, A., Al-Kahtani, H. (1993). Composition of Ghee (Samn Barri's) from Cow's and Sheep's Milk. *Food Chem*, 46, 373-375.



# STARDOM UNIVERSITY

Stardom Journal of Engineering and Natural Studies



— STARDOM JOURNAL OF ENGINEERING AND NATURAL SCIENCES —

PUBLISHED TWICE A YEAR BY STARDOM UNIVERSITY

Volume 2 - 1<sup>st</sup> issue 2024

International deposit number : ISSN 2980-3756