

Synthesis, Identification and Assess the Biological and Laser Efficacy of New Compounds of Azetidine Derived from Benzidine

Adil Hussein Dalaf¹ and Fawzi Hameed Jumaa²

¹Department of Chemistry - College of Science - Tikrit University. Tikrit-Iraq

²Department of Chemistry - College of Education for Women - Tikrit University. Tikrit-Iraq

*Corresponding Author: adil.h.dalaf@tu.edu.iq

Received 23/9/2020 , Accepted 11/10/2020 , published:14/10/2020

DOI: 10.18081/2226-3284/14-10/12-25

Abstract: In this study, new azetidine [A₁-A₅] compounds were prepared by the reaction five of schiff bases compounds with chloroacetyl chloride in 1,4-dioxane. The prepared compounds were characterized by physical properties, UV-Vis, ¹H-NMR and FT-IR spectral and C.H.N analysis. TLC checked the purity for these compounds. The antibacterial activities were studied against different kinds of bacteria, namely *Staphylococcus epidermidis* and *Staphylococcus aureus* Gram (+) ve, *Escherhshia coli*, and *Klebsiella Pneumonia* Gram (-) ve. Also, evaluation of laser efficacy is shown for the compounds (A₁-A₅) were laser radiated to (10, 20, 30) seconds. When the melting point and color were measured, it was found that the compounds were not affected or polymerized.

© 2020 Al Muthanna University. All rights reserved.

Keywords: Benzidine, Azetidine, Biological activity, Laser Effectiveness.

1. Introduction

Azetidines are present in various natural products (amino acids) [1], and studies have shown that integrating azetidines into pharmaceutically applicable scaffolds can lead to improved pharmacokinetic properties and metabolic stability [2,3]. Methods developed for azetidine synthesis depend on nucleophilic replacement reactions with nitrogen nucleophiles, reduction of β -lactams [4], or ring opening of highly strained azabicyclobutanes [5]. Considering that most of these approaches require functionalized starting materials and involve multistep sequences, aza Paternò-Büchi reactions represent a highly efficient and direct strategy

for the synthesis of azetidines [6]. The azetidine group is one of the most important pharmaceutical substructures and is present in a variety of approved drugs such as antibiotics [7] kinases (e.g. baricitinib, cobimetinib) [8], and other groups of compounds (e.g. thrombin inhibitor, azelnidipine) [9]. In addition, the azetidine group reduces the toxicity of the raw materials prepared from it [10] and lead to compounds with an enhanced pharmacokinetic [11]. The enhanced interest in azetidines needs new synthetic methods with varying substitution and stereochemistry to reach desired scaffolds [12]. Substituted azetidines strongly attract chemists because of their importance in catalysis, stereoselective

synthesis and medicinal chemistry [13]. Other techniques achieve the creation of rings and stereocenters simultaneously by uniting two fragments; however, this also leads to difficulties in determining the proper regio- and stereochemistry [14]. The reactions (2+2) were well-explored [15] while the related reactions (3+1) are less common [16].

2. Experimental

2.1. Materials and devices: All chemicals (Chloroacetyl chloride, triethylamine, 1,4-dioxane and dimethyl sulfoxide) had been used were supplied by company Alfa Aesar. The melting points were determined by Electro thermal Melting Apparatus 9300 in open capillary tubes that were uncorrected. Thin-layer chromatography (TLC) was used to track and verify the purity of the reaction. The Fourier-transform infrared spectroscopy (FT-IR) spectra were recorded using FT-IR 8400S Shimadzu Scale (4000-400) cm⁻¹ spectrophotometer. In addition, the UV-Vis. Spectra in ethanol were measured using Shimadzu 800UV in range (200-400) nm. Proton Nuclear Magnetic Resonance (¹H-NMR) Spectra was recorded on Varian operating at 400 MHz instrument using dimethyl sulfoxide (DMSO-d⁶) as a solvent. Quantitative analysis of the spectrophotometer elements determined using elements analysis (C.H.N). The prepared compounds were irradiated with helium-neon laser beam

(visible laser) of 1 milliwatt and wavelength 600-700 nm, 2010 model.

2.2. Synthesis methods

2.2.1. Synthesis of azetidine [A₁–A₅]

Mixture of Schiff bases [17] (0.001 mol) with Et₃N (0.004 mol, 0.55 ml) in (10 ml) 1,4-dioxane then chloroacetyl chloride (0.004 mol, 0.3 ml) was added dropwise for a period slowly of 30 min. The reaction mixture was refluxed for (9-12) hrs. So when the reaction was terminated- (TLC identified), the mixture was placed into ice-cold water for solid residue output. Filtering, drying, and recrystallizing of benzene and ether to the effects (50-50) [18]. Physical properties are given in table (1).

2.3. Antibacterial Activity

The measurement of antibacterial activity of compounds [A₁-A₅] against gram positive and negative such as *Escherchia coli*, *Klebsiella Pneumonia* are Gram (-) ve and *Staphylococcus epidermidis*, *Staphylococcus aureus* are Gram (+) ve by using the disk diffusion method. They sprayed the disks with a DMSO. Then, incubate them overnight at 37 °C. The negative omission was DMSO. The maximal inhibition zone against the development form of test microorganism was observed and calculated for analysis. *Ampicillin*, *amoxicillin*, and *Ciprofloxacin* were used as control samples at three concentrations [19, 20].

3. Results and Discussion

In this research, five compounds were prepared including azetidine [A₁-A₅] as in the scheme (1) and Identification by UV-Vis, ¹H-NMR, FT-IR Spectra and C.H.N analysis.

3.1. Characterization of azetidine [A₁-A₅]

The azetidine [A₁-A₄] were prepared by the reaction (1 mole) schiff base with (2 mole) of chloroacetyl chloride in 1,4-dioxane.

The UV spectra showed the transitions $n \rightarrow \pi^*$ and $\pi \rightarrow \pi^*$ indicated the existence of unbonded pair electrons on nitrogen, oxygen atoms, and aromatic (double bond) framework.

The FT-IR spectrum showed azetidine disappearance of band azomethine (C=N) group (1622-1672) cm⁻¹, beside new bands appear at (2906-2988) cm⁻¹ attributed to the (CH) aliphatic as well as the emergence of band at (1650-1666) cm⁻¹ due to (C=O) for azetidine. Besides other bands at (3369-33463) cm⁻¹ for (OH) and at (3033-3082) cm⁻¹ for aromatic (C-H), in addition, at (1573-1598) cm⁻¹ and at (1469-1498) cm⁻¹ due to (C=C) aromatic ring, and at (1421-1444) cm⁻¹ for azo (N=N), and at (840-880) cm⁻¹ for (C-Cl), as shown in fig. (1,2,3). U.V and FT-IR spectrum are given in table (2) [21].

In addition, the ¹H-NMR spectrum of [A₁] fig. (4) displays simple single signal at $\delta = 9.40$

ppm attributed to (OH) (A) and a multiple signal in the range $\delta = (9.05-7.10)$ ppm for the aromatic protons (B,C,D,E,F), and doublet signal at $\delta = (6.47 \ \& \ 6.45)$ ppm attributed to (CH-Cl) (G), and doublet signal at $\delta = (6.09 \ \& \ 6.07)$ ppm attributed to (CH-N) (H) in addition to singlet signal at $\delta = 2.50$ ppm attributed to DMSO-d⁶ (I).

In addition, the ¹H-NMR spectrum of [A₃] fig. (5) displays simple singlet signal at $\delta = 10.15$ ppm attributed to (OH) (A) and doublet signal at $\delta = (9.23 \ \& \ 9.24)$ ppm attributed to (N=CH) (B) in the ring and a multiple signal in the range $\delta = (8.51-7.85)$ ppm for the aromatic protons (C,D,E,) and doublet signal at $\delta = (7.50 \ \& \ 7.48)$ ppm attributed to (CH) (F) and doublet signal at $\delta = (6.19 \ \& \ 6.18)$ ppm attributed to (CH-Cl) (G), and doublet signal at $\delta = (5.99 \ \& \ 5.95)$ ppm attributed to (CH-N) (H) as well as singlet signal at $\delta = 2.50$ ppm attributed to DMSO-d⁶ (I). [22-24].

3.2. Antibacterial activity

The effect of the prepared compounds [A₁-A₅] on the growth of bacteria, namely *Escherchia coli*, *Klebsiella Pneumonia Gram (-ve)*, *Staphylococcus epidermidis* and *Staphylococcus aureus Gram (+ve)* was determined. Antibacterial activity of the prepared compounds were studied and the results showed that some of the prepared compounds possess good antibacterial activity. The results of inhibition zone (IZD) in

millimeters are shown in table (3) [25], see fig. (6,7) and see scheme (2–5).

3.3. Influence of lasers on prepared compounds [A₁-A₅]

A laser system with a power of (5) milliwatt that gives laser rays in continuous waves in the visible region of the spectrum at a wavelength (600-700) nm. Laser radiation of compounds [A₁-A₅] lasered for (10, 20, 30) seconds. It has been found the compounds have not been

affected. When the melting point and color are determined, they did not disintegrate or polymerize. This denotes that the laser beams used did not affect the compounds. Since they are stable, as shown in the table (4) [26].

Table (1): Physical properties and elemental analysis of prepared compounds [A₁-A₅]






Comp. No.	Ar	Molecular Formula M. Wt.	Color	M.P (C ⁰)	T. hr.	Y. %	R.f. MeOH	found / (calc.) %		
								C%	H%	N%
A 1		C ₄₂ H ₃₀ C ₁₂ N ₆ O ₄ 753.64	Brown	199-200	11	73	0.90	66.85 (66.94)	4.07 (4.01)	11.07 (11.15)
A 2		C ₄₂ H ₂₈ C ₁₂ N ₈ O ₈ 843.63	Black	164-166	12	81	0.72	59.71 (59.80)	3.36 (3.35)	13.24 (13.28)
A 3		C ₃₈ H ₂₆ C ₁₂ N ₁₀ O ₄ 757.59	Brown	257-258	9	72	0.88	60.20 (60.25)	3.51 (3.46)	18.42 (18.49)
A 4		C ₄₂ H ₃₀ C ₁₂ N ₆ O ₆ 785.64	Dark brown	237-239	9	43	0.68	64.15 (64.21)	3.89 (3.85)	10.66 (10.70)
A 5		C ₅₀ H ₃₈ C ₁₂ N ₁₀ O ₁₀ S ₂ 1073.93	Gray	210-212	12	57	0.73	55.90 (55.92)	3.61 (3.57)	12.98 (13.04)

Table (2): FT-IR and UV/Vis data of prepared compounds [A₁-A₅]






Comp. No.	Ar	λ ₁ max nm λ ₂ max nm	IR (KBr) cm ⁻¹						
			ν O-H	ν C-H Arom.	ν C-H Aliph.	ν C=O	ν C=C	ν N=N	ν C-Cl
A 1		227 394	3463	3072	2941	1652	1579 1483	1421	840
A 2		246 395	3369	3033	2925	1656	1598 1496	1444	858
A 3		257 336	3401	3064	2906	1650	1587 1875	1439	871
A 4		263 397	3386	3058	2920	1666	1573 1465	1429	854
A 5		206 383	3426	3082	2988	1659	1591 1498	1438	880


Table (3): Antibacterial activity of the prepared compounds [A₁-A₅] and control antibiotic

Comp. No.	<i>E. Coil</i>	<i>K. Pneumonia</i>	<i>S. Aureus</i>	<i>S. Epidermidis</i>
-----------	----------------	---------------------	------------------	-----------------------

	Conc. mg/ml			Conc. mg/ml			Conc. mg/ml			Conc. mg/ml		
	25	50	100	25	50	100	25	50	100	25	50	100
A ₁	2	3	4	1	3	5	2	4	5	0	2	3
A ₂	2	3	5	2	2	4	0	1	2	1	3	4
A ₃	1	3	4	2	4	4	1	2	3	1	2	3
A ₄	0	2	3	2	3	5	0	1	2	1	3	4
A ₅	3	5	5	2	3	4	2	4	5	2	3	4
<i>Amoxicillin</i>	2	3	4	2	4	4	2	3	4	1	2	3
<i>Ampicillin</i>	2	4	4	2	3	3	2	3	4	2	2	3
<i>Ciprofloxacin</i>	2	3	3	2	3	4	1	2	3	1	3	4
Blank disk	0	0	0	0	0	0	0	0	0	0	0	0

Table (4): The results of the irradiation of the compounds by laser beams

Comp. No.	10 S		20 S		30 S	
	M.P. °C	Color	M.P. °C	Color	M.P. °C	Color
A ₁	199-200	Brown	199-200	Brown	199-200	Brown
A ₂	164-166	Black	164-166	Black	164-166	Black
A ₃	257- 258	Brown	257- 258	Brown	257- 258	Brown
A ₄	237-239	Dark brown	237-239	Dark brown	237-239	Dark brown
A ₅	210-212	Gray	210-212	Gray	210-212	Gray



Scheme (1): Route of prepared compounds [A₁-A₅]




Fig. (1): FT-IR spectrum of compound [A₁]




Fig. (2): FT-IR spectrum of compound [A₂]




Fig. (3): FT-IR spectrum of compound [A₄]




Fig. (4): ¹H-NMR spectrum of compound [A₁]




Fig. (5): ¹H-NMR spectrum of compound [A₃]




Fig. (6): Antibacterial activity of compound [A₁] against *E. coli*.







Fig. (7): Antibacterial activity of compound [A₅] against *K. pneumoniae*.




Scheme (2): Evaluation of inhibitory activity of compounds prepared for *Escherichia Coli*



Scheme (3): Evaluation of inhibitory activity of compounds prepared for *K. Pneumonia*



Scheme (4): Evaluation of inhibitory activity of compounds prepared for *S. Aureus*



Scheme (5): Evaluation of inhibitory activity of compounds prepared for *S. Epidermidis*

4. Conclusions

The spectroscopic measurements showed the purity of the compounds prepared. In addition, some of the prepared compounds showed good antibacterial activity against the bacterial such as *Eschershia coli*, *Klebislla Pneumonia Gram (-ve)*, *Staphylococcus epidermidis* and *Staphylococcus aureus Gram (+ve)*. Laser radiation of compounds [A₁-A₅] for (10, 20, 30) seconds was studied. It has been found the compounds have not been affected. When the melting point and color were determined, they neither disintegrated nor polymerized. This clarified that the laser beams used did not affect the stable compounds.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

The authors are grateful for everyone who supported us in getting the chemicals.

The authors also thank for everyone who helped us calculate, analyze and interpret the results.

Credit authorship contribution statement

Adil H. Dalaf: Conceptualization, Methodology, Investigation, Software, Validation, Data curation.

Fawzi H. Jumaa: Supervision, Investigation, Writing - review & editing.

References

[1] D'Auria, M. (2019). The Paternò–Büchi reaction–a comprehensive review. *Photochemical & Photobiological Sciences*, 18(10), 2297-2362.

- [2] Majumdar, K. C., & Chattopadhyay, S. K. (Eds.). (2011). *Heterocycles in natural product synthesis*. John Wiley & Sons.
- [3] Brown, A., Brown, T. B., Calabrese, A., Ellis, D., Puhalo, N., Ralph, M., & Watson, L. (2010). Triazole oxytocin antagonists: identification of an aryloxyazetidone replacement for a biaryl substituent. *Bioorganic & medicinal chemistry letters*, 20(2), 516-520.
- [4] Berthe, B., Outurquin, F., & Paulmier, C. (1997). Synthesis of azetidines by electrophilic selenium-induced cyclization of homoallylic benzylamines. *Tetrahedron letters*, 38(8), 1393-1396.
- [5] Gianatassio, R., Lopchuk, J. M., Wang, J., Pan, C. M., Malins, L. R., Prieto, L., & Spangler, J. E. (2016). Strain-release amination. *Science*, 351(6270), 241-246.
- [6] Freneau, M., & Hoffmann, N. (2017). The Paternò-Büchi reaction-Mechanisms and application to organic synthesis. *Journal of Photochemistry and Photobiology C: Photochemistry Reviews*, 33, 83-108.
- [7] Reidl, T. W., Son, J., Wink, D. J., & Anderson, L. L. (2017). Facile synthesis of azetidine nitrones and diastereoselective conversion into densely substituted azetidines. *Angewandte Chemie International Edition*, 56(38), 11579-11583.
- [8] Rice, K. D., Aay, N., Anand, N. K., Blazey, C. M., Bowles, O. J., Bussenius, J. & Engst, S. (2012). Novel carboxamide-based allosteric MEK inhibitors: discovery and optimization efforts toward XL518 (GDC-0973). *ACS medicinal chemistry letters*, 3(5), 416-421.
- [9] Sorbera, L. A., Bayes, M., Castaner, J., & Silvestre, J. (2001). Melagatran and ximelagatran. *Drug Future*, 26, 1155-1170.
- [10] Yan, Q., Wang, Y., Zhang, W., & Li, Y. (2016). Novel Azetidine-Containing TZT-1027 Analogues as Antitumor Agents. *Marine drugs*, 14(5), 85.
- [11] Kirichok, A. A., Shton, I. O., Pishel, I. M., Zozulya, S. A., Borysko, P. O., Kubyskin, V. & Mykhailiuk, P. K. (2018). Synthesis of multifunctional spirocyclic azetidines and their application in drug discovery. *Chemistry—A European Journal*, 24(21), 5444-5449.
- [12] Schmid, S. C., Guzei, I. A., & Schomaker, J. M. (2017). A stereoselective [3+ 1] ring expansion for the synthesis of highly substituted methylene azetidines. *Angewandte Chemie*, 129(40), 12397-12401.
- [13] Antermite, D., Degennaro, L., & Luisi, R. (2017). Recent advances in the chemistry of metallated azetidines. *Organic & biomolecular chemistry*, 15(1), 34-50.
- [14] Sakamoto, R., Inada, T., Sakurai, S., & Maruoka, K. (2016). [2+ 2] Photocycloadditions between the carbon-

nitrogen double bonds of imines and carbon–carbon double bonds. *Organic letters*, 18(24), 6252-6255.

[15] Ghosh, A., Mandal, S., Chattaraj, P. K., & Banerjee, P. (2016). Ring expansion of donor–acceptor cyclopropane via substituent controlled selective N-transfer of oxaziridine: synthetic and mechanistic insights. *Organic letters*, 18(19), 4940-4943.

[16] Han, J. Q., Zhang, H. H., Xu, P. F., & Luo, Y. C. (2016). Lewis acid and (hypo) iodite relay catalysis allows a strategy for the synthesis of polysubstituted azetidines and tetrahydroquinolines. *Organic letters*, 18(20), 5212-5215.

[17] Dalaf, A. H., Jumaa, F. H., & Jabbar, S. A. S. (2018). Synthesis and Characterization of some 2, 3-dihydroquinoxaline and evaluation of their biological activity. *Tikrit Journal of Pure Science*, **23(8)**: 66-67.

[18] Salwa, A. J., Ali, L. H., Adil, H. D., Hossam, S. A. (2020). Synthesis and Characterization of Azetidine and Oxazepine Compounds Using Ethyl-4-((4-Bromo Benzyldiene) Amino) Benzoate as Precursor and Evaluation of Their Biological Activity. *Journal of Education and Scientific Studies*, ISSN: 24134732. **16(5)**: 39-52.

[19] Abd, I. Q., Ibrahim, H. I., Jirjes, H. M., & Dalaf, A. H. (2020). Synthesis and Identification of new compounds have Antioxidant activity Beta-carotene, from

Natural Auxin Phenyl Acetic Acid. *Research Journal of Pharmacy and Technology*, **13(1)**: 40-46.

[20] Saleh, R. H., Rashid, W. M., Dalaf, A. H., Al-Badrany, K. A., & Mohammed, O. A. (2020). Synthesis of Some New Thiazolidinone Compounds Derived from Schiff Bases Compounds and Evaluation of Their Laser and Biological Efficacy. *Ann Trop & Public Health*, **23(7)**: 1012-1031.

[21] PATEL, S., SEN, A. K., DASH, D. K., SADHU, P., & BAILE, S. B. (2020). Synthesis, Characterization and Biological Evaluation of Azetidine Analogues and Related Compounds. *International Journal of Pharmaceutical Research*, 12(2).

[22] Rani, V. E., & Reddy, P. R. (2018). Synthesis and Antimicrobial Activity of New Pyridine Containing Substituted Phenyl Azetidine-2-One Derivatives. *Open Journal of Medicinal Chemistry*, 8(2), 22-29.

[23] Dhall, E., Jain, S., Mishra, A., Dwivedi, J., & Sharma, S. (2018). Synthesis and Evaluation of Some Phenyl Substituted Azetidine Containing 1,2,4-triazole Derivatives as Antibacterial Agents. *Journal of Heterocyclic Chemistry*, 55(12), 2859-2869.

[24] Hasan, S. M., Majeed, I. Y., & Samir, A. H. (2018). Synthesis and Characterization of Some New Dioxoisindolin Compounds Containing Thiazine, Azetidine, Thiazolidine

and Amide Moities. Diyala Journal For Pure Science, 14(03), 94-107.

[25] Ivanišová, E., Frančáková, H., Ritschlová, P., Dráb, Š., Solgajová, M., & Tokár, M. (2020). Biological activity of apple juice enriched by herbal extracts. Journal of Microbiology, Biotechnology and Food Sciences, 9(4): 69-73.

[26] Salih, B. D., Dalaf, A. H., Alheety, M. A., Rashed, W. M., & Abdullah, I. Q. (2020). Biological activity and laser efficacy of new Co (II), Ni (II), Cu (II), Mn (II) and Zn (II) complexes with phthalic anhydride. Materials Today: Proceedings. 1-6.