

Monthly variations of some physical and chemical properties for Al-Garraaf River one of the main Tigris branches at Al-Haay Cit

Wisam Th. Al-Mayah and Muhammad N. Al-Azzawi

University of Baghdad / College of Science / Biology Department

Iraq – Baghdad

Abstract

Monthly variations in Physio-chemical parameters of Al-Garraaf, one of the main tributaries of the Tigris river were investigated from October 2012 to April 2013. The study location situated in the south-eastern sector of Iraq and surrounded by wide and fertile agricultural lands. Three stations were selected to. The former is located at 2 km of AL-Haay City as control. The second is situated at distance 2 km away from the former represented study area and the latter station is located at 4 km apart from the second one. collecting samples monthly ,two samples were taken each month .

In the present study Fourteen physical and chemical parameters were analyzed based on the importance of these parameters. These Fourteen parameters are ranged as following: air temperature (16 to 32) °C, water temperature (11 to 26) °C, pH (7.05to 8.35), E.C. (827 to1558) $\mu\text{S}/\text{cm}$, salinity (0.54 to 0.99) ppt, DO (7.04 to 10.38) mg/L , BOD (0.90 to 7.01) mg/L ,turbidity (31.0 to 177.0) NTU, TDS (550.78 to 1108) mg/L , TSS (38 to 636.5) mg/L , T.H (275 to 500) mg/L , Cl^- (89 to 184.6) mg/L, NO_3 (5.7 to 15.76) mg/L and PO_4 (0.15 to 0.5) mg/L.

1. Introduction

Rivers have always been the most important fresh water resources, and most developmental activities are still dependent upon them. Rivers play a major role in assimilating or carrying industrial and municipal waste water, manure discharge and runoff which are responsible for water river pollution[1] [2]

The degree of pollution is generally assessed by studying physical and chemical characteristics of the water bodies[3]

Iraqi inland waters witness tremendous impacts through discharges of manufacturers, agricultural and domestic sewage [4] [5]. Quite few studies were performed on Tigris River [6] [7] [8], but no work had considered Al-Garraf canal in Al-Haay City. The present study has taken in consideration the investigation of abiotic conditions in this vital habitat on monthly basis .

2. Material and Methods

2.1. Study area

Al- Garraf River is one of two branches of the Tigris River at Kutt City, 225 km south of Baghdad City (Fig. 1). After branching from the Tigris, the Garaff flows southeast toward Al-Haay City (study area) within Wasit Province, 220 km southwest of Baghdad City. The river is 230 km in length with a variable depth of 17.4 m at its branching point from the Tigris to 10.0 m at its junction with the Euphrates River at the marsh area near Thiqrar Province .



Figure (1) showing map of sampling location in the study area

2.2.Sampling

Samples for physical and chemical variables were performed from three sites during period extended from the October 2012 to April 2013. Water samples were collected for physiochemical analysis using pre-washed polyethylene bottle by water sample twice before filling.

The studied physico-chemical parameters include water temperature (by using precise mercury thermometer), hydrogen ion concentration (by using pH-meter), electrical conductivity (by using EC-meter), turbidity level (by using turbidity-meter), dissolved oxygen (titrimetric methods), biological oxygen demand (Winkler methods), nitrate, reactive phosphate (by using spectrophotometric methods), total hardness and chloride (by using titrimetric methods), were measured according to APHA[9] [10].

3.Results

Figure (2) and Table (1), shows monthly changes in air temperature for the three selected stations. Values ranged between 16°C in station-1 during January (2013) to 32° C in station-2 during April (2013).

Figure (3) and Table (1), however, indicates monthly variations in water temperature. The lowest value was 11°C in station-2 during January (2013) and the highest 26°C in station-2 during April (2013).

Figure (4) and Table (1), shows monthly changes in pH. The lowest (7.05) was encountered in October (2012) from station-2 and the highest (8.35) was recorded in November (2012), but values in general were slightly alkaline direction.

Figure (5) and Table (1), Shows monthly changes in values of Electrical conductivity. The lowest (827 $\mu\text{s}/\text{cm}$) was measured from station-3 in March (2013) and the highest (1558 $\mu\text{s}/\text{cm}$) was observed in October (2012) from station-2.

Figure (6) and Table (1), shows monthly changes in values of water salinity. The lowest (0.54 ppt) was observed in station-3 during March (2013) and the highest (0.99 ppt) measured from station-2 in October (2012). Figure (7) and Table (1), Shows monthly changes in values of Turbidity. The lowest (31.0 NTU) was

observed in station-3 in December (2012) and the highest (177.0 NTU) was observed in January (2013) from station-2.

Figure (8) and Table (1), Reveals monthly variations in dissolved oxygen in selected stations Values declined during October (2012). The lowest (7.04 mg/L) was in October (2012) from station-2 and the highest (10.38 mg/L) was, in general, in January (2013) from station-1.

Figure (9) and Table (1), shows monthly variations in values of biological oxygen demands (BOD). The lowest (0.90 mg/L) was recorded in March (2013) from station-1 and the highest (7.01 mg/L) was in November (2012) from station-2.

Figure (10) and Table (1), shows monthly changes in total dissolved solid. The lowest (550.78 mg/L) was encountered in March (2013) from station-3 and the highest (1108 mg/L) was recorded in October (2012) from station-2.

Figure (11) and Table (1), shows monthly variations in total suspended solid. The lowest (38 mg/L) was observed in December (2012) from station-1 and the highest (636.5 mg/L) was observed in January (2013) from station-2.

Figure (12) and Table (1), Revealed monthly variations in values of total hardness in the selected localities. Highest value (500 mg/L) was in November (2012) and encountered from station-2. The lowest (275 mg/L), however, was in January (2013) from station-1.

Figure (13) and Table (1), Shows monthly changes in values of Chloride. The lowest (89.02 mg/L) was measured from station-1 in March (2013) and the highest (184.6 mg/L) was observed in November (2012) from station-2.

Figure (14) and Table (1), Reveals monthly variations in Nitrate. The lowest (5.7 mg/L) was in April (2013) from station-1 and the highest (15.76 mg/L) was observed in November (2012) from station-2.

Figure (15) and Table (1), Shows monthly changes in values of Reactive phosphate. The lowest (0.15 mg/L) was observed in October (2012) from station-1 and the highest (0.5 mg/L) was observed in December (2012) from station-2.

Table (1): Monthly Variation for Al-Gharaf River through Period study 2012 – 2013


Time	Station	Air T. C °	Water T. C °	PH	E.C ms/cm	Sali. mg/l	DO mg/l	DOB5 mg/l
October 2012	Station1	30	24	7.38	1335	0.87	8.22	1.89
	Station2	30	26	7.05	1558	1.0	7.04	7.0
	Station3	32	25	7.34	1339	0.87	7.61	4.01
November 2012	Station1	25	20	7.7	910	0.59	9.62	1.45
	Station2	25	17	7.06	1150	0.75	8.44	7.01
	Station3	24	20	8.35	915	0.59	8.25	3.14
December 2012	Station1	23	19	8.1	1228	0.79	8.01	2.08
	Station2	22	17	8.1	1387	0.90	8.25	6.39
	Station3	23	17.5	8.2	1311	0.86	8.59	4.12
January 2013	Station1	16	11.5	8.09	1075	0.69	10.38	2.0
	Station2	16	11	7.98	1078	0.70	8.36	5.5
	Station3	16	12.5	7.77	1062.5	0.69	9.57	2.5
March 2013	Station1	28	24	7.8	855	0.56	8.0	0.90
	Station2	29	23	7.5	888	0.58	7.5	4.5
	Station3	30	23	7.8	827	0.54	8.1	2.1
April 2013	Station1	30	25	7.8	1018	0.65	8.1	1.9
	Station2	32	26	7.3	1029	0.66	7.2	5.2
	Station3	31	24	7.2	1014	0.65	7.9	2.9

Appendix table-1


Time	Station	Turb. NTU	TSS mg/l	TDS mg/l	T.H mg/l	Cl⁻ mg/l	NO3⁼ mg/l	PO4⁻³ mg/l
October 2012	Station1	78	838	946	430	99.4	15.09	0.15
	Station2	121	1090	1108	460	134.9	15.00	0.57
	Station3	74	938	928	437.5	113.6	15.69	0.31
November 2012	Station1	33.0	828	606.1	360	118.9	6.75	0.17
	Station2	56.0	1132	765.9	500	184.6	15.76	0.19
	Station3	53.0	834	609.2	360	124.3	7.21	0.23
December 2012	Station1	33.0	38	817.9	407	133.0	6.6	0.3
	Station2	42.0	74	923.8	438	152.0	7.0	0.5
	Station3	31.0	85	873.2	415	139.0	7.5	0.4
January 2013	Station1	161.0	976	715.9	275	134.9	10.12	0.19
	Station2	177.0	986	717.9	335	138.5	12.45	0.23
	Station3	163.5	948	707.7	360	120.7	9.62	0.2
March 2013	Station1	35	65	569.43	310.4	89.02	6.1	0.32
	Station2	67	96	591.40	388	103.07	7.0	0.43
	Station3	43	85	550.78	329.8	91.36	6.0	0.38
April 2013	Station1	33	56	677.99	314	89	5.7	0.33
	Station2	52	86	685.32	412	109	6.2	0.45
	Station3	37	87	675.32	392	95	5.7	0.38

Table (2): Comparison between some water quality parameters of Al-Gharraf River with the Iraqi and international standards.


Parameter	WHO standards for drinking water in 2004	EEC 464/76 standard for surface water quality (Tebbutt, 1998)	Iraqi standards for water quality in 1998	Present Study Mean
pH	6.5- 8.5	6.5- 8.5	6.5- 8.5	7.7
TDS mg/L	500-1500	500-1500	1000	829.39
Turbidity NTU	0-50	0-25	25	104
DO mg/L	> 5	> 5	> 5	8.71
BOD5 mg/L	> 3	> 5	> 3	3.955
T.H mg/L	100 - 500	100-500	-	387.5
Cl ⁻ mg/L	200	200 – 600	200	136.81
NO ₃ ⁻ mg/L	0 – 45	0 – 40	25 –50	10.73
PO ₄ mg/L	0.1	0.4	0.4	0.325




(Fig.2) Monthly changes in Air Temperature in three selected stations




(Fig.3) Monthly changes in Water Temperature in three selected stations




(Fig.4) Monthly changes in Hydrogen concentrated in three selected stations




(Fig.5) Monthly changes in Electrical Conductivity in three selected stations




(Fig.6) Monthly changes in salinity concentrated in three selected stations




(Fig.7) Monthly changes in Turbidity in three selected stations




(Fig.8) Monthly changes in Dissolved Oxygen in three selected stations




(Fig.9) Monthly changes in Biological oxygen demand in three selected stations




(Fig.10) Monthly changes in Total Dissolved solids in three selected stations




(Fig.11) Monthly changes in Total Suspended solids in three selected stations




(Fig.12) Monthly changes in Total Hardness in three selected stations



(Fig.13) Monthly changes in Chloride in three selected stations



(Fig.14) Monthly changes in Nitrate concentrated in three selected stations



(Fig.15) Monthly changes in Phosphate concentrated in three selected stations

4. Discussion

Air and water temperature is an important factor in any aquatic environments affecting on biological processes, in this study it was ranged between 16 to 32 °C and 11 to 26 °C respectively. This result was similar to previous studies done by [11] [12].

The pH value of AL-Garraf River in study sites during of most studied period was alkaline side above 7, and this result agreed with [13], they reported that Iraqi inland water is regarded to be on the alkaline side of neutrality, reflecting geological formations of the area and the results are agree with the finding that recorded by [14] [15].

Electrical conductivity used as an indicator of water quality based on total dissolved salts [16]. The increase EC values at station two reflects the strong effect of domestic sewage effluent discharge at this area. Also, EC values recorded in the present work is coincided with findings of [17] [18].

The study also revealed monthly changes in salinity, with notable increase during summer months due to evaporation [19]. The presence of agricultural drainage systems namely, Kut, Al-Muafakiah and AL-Haay may contribute in rising salinity as well.

Water turbidity is caused by suspended matter such as clay, silt and planktons also turbidity degree of River water is an approximate measure of the intensity of the pollution [20]. This result was similar to previous studies done by [21] [22].

Oxygen content of water is one of the important factors, and it is very necessary for all living organisms [23]. The study finding coincided with other authors [24] [25] [26] on Iraqi inland waters mainly Tigris. Low concentration of DO recorded from station-2 may relate to organic wastes discharged from AL-Haay City. Generally, the DO at most stations of canal water was within normal guideline values cited by [27] for the protection of aquatic life.

The biological oxygen demand is defined as the quantity of DO which is able to oxidize the organic components in the water with the assistance of microorganisms under defined experimental conditions [28].

Generally, results indicate increasing levels of BOD, in particular at station-2 during the November and October, this may be due to decomposition of organic matters run directly to the river with domestic sewage. These results were slightly higher than that reported by [29] [30] at the same river.

Values of total hardness in the selected stations exceeded 490 mg/L as CaCO₃. This indicates that waters are very hard according to [31]. Increase in hardness values was found to coincide with rise in salinity [32] [33]. The results of total hardness are agreed with those of [34] [35].

Chloride is a natural substance present in all portable water as well as sewage effluents as metallic salt. Generally high concentration of chloride indicates to organic pollution in the water [36]. This result was similar to previous studies done by [37] [38].

Total dissolved solids (TDS) is the term used to describe the inorganic salts and small amounts of organic matter present in solution in water [39]. These results were slightly lower than that reported by [40] [41] at the same river.

Total suspended solids are considered to be one of the major pollutants that contributes to the deterioration of water quality, contributing to higher costs for water treatment, decreases in fish resources, and the general aesthetics of the water [42], TSS values recorded in the present work is coincided with findings of [43] [44].

Nitrate is the stable form of combined nitrogen and it is an important factor which might limit growth of phytoplankton [45]. The results of nitrate are agreed with those of [46] [47].

Phosphorus is essential to the growth of algae and other biological organisms. The reactive phosphate concentration in studied river was ranged between 0.1 to 0.5 mg/l. The high concentration of phosphate may be due to sewage water effluent and fertilizer application in surrounding agricultural area. This result was close to that reported by [48] [49].

5- Conclusions

- 1- The waste of factories, urban run-off, city sewage and the agricultural activities were affecting the physicochemical characteristics of Al-Garraf River.
- 2- The results revealed that water parameters were most within the Iraqi standards, and WHO standards for the raw water .

6- References

- 1- **Toman**, M.J.(2009). Physico - Chemical Characteristics and Seasonal Changes of plankton communities in a reservoir. *Lakes and reservoirs research and management*. 2(1&2); 71 - 76.
- 2- **Kumar**, Rita. N., Rajal Solanki and Nirmal Kumar J.I (2011). An Assessment of Seasonal Variation and Water Quality Index of Sabarmati River and Kharicut Canal at Ahmedabad, Gujarat *Electronic Journal of Environment, Agriculture and Food Chemistry* 10 (8), 2771-2782
- 3- **Duran**, Mustafa and Menderes Suicmez (2007). Utilization of both benthic macroinvertebrates and physiochemical parameters for evaluating ;water quality of the stream Cekerek (Tokat Turkey).*J. Environ. Biol* 28,231-236.
- 4- **Wrobel**, L. C. and Brebbia, C. A. (1994). Modeling measuring and production in : water pollution Wessex Institute of technology. U. K. PP.51.
- 5- **Klerks**, P. L. and Lentz, S. A. (1998). Resistance to lead and zinc in the western Mosquito fish, *Gambusia affinis* Inhabiting contaminated bayon tre pagnier, *Ecotoxicology*, 7 : 11-17.
- 6- **Al-Lami**, A. A. ; and Th. I. Kassim ; A. W. Sabri and K. A. Rasheed (1996). The ecological effects of Diyala river on Tigris River. I. *Limnology. J. Coll. Educ. for women. Univ. Baghdad*, 7(1):84-92.
- 7- **Al-Saadi**, H. A.; Al-Lami, A. A.; Kassim, T. I. (1999) studies to ecological characteristics of upper Tigris and Euphrates Rivers and its correlation with fishery progress in Iraq. *J. Environmental and sustainable progress researches*, Vol.2, No.2- 1420. p. 24-31. (in Arabic)

- 8- **Al-Rubae**, M. A. J. (1997) Ecological study to Ethem river and its impact on Tigris. MSc thesis; Baghdad University. (in Arabic)
- 9- **APHA**(1998). Standard methods for the examination of water and waste water, 20th ed.
- 10- **APHA**,AWWA and WFF (2005) Standard Methods for the Examination of Water and wastewater, 21th ed., edited by Eaton, A. D. ; L. S. Clesceri; E. W. Rice, and A. E. Greenberg. American Water Work Association and Water Environment Federation,USA.
- 11- **Sabri**, A. W. ; Maulood, B. K. and Sulaiman, N. E. (1989). Limnological studies on river Tigris : Some physical and chemical characters. J. Biol. Sci. Res., 20 (3) : 565-579.
- 12- **Al-Rubae**, M. A. J. (1997) Ecological study to Ethem river and its impact on Tigris. MSc thesis; Baghdad University. (in Arabic)
- 13- **Maulood** B K, Hinton G C F, and Al-Dosky H S. 1980. A study on the blue green algal flora of Arbil province, Iraq. Zanco Sci. J. Univ. Sulaimaniyah, Iraq, 6:67-90.
- 14- **Al-Lami**,A.A.; Jaweir, H. J. and Nashaat, M. R. (1998). Benthic invertebrates of the river Euphrates upstream and downstream sectors of Al-Qadisia Dam, Iraq. Reg. Riv. 14 (4) : 383-390 .
- 15- **Al-Obaidi**, G. S. A. (2006). A Study of phytoplankton community in Abu Zirig marsh, Southern Iraq. MSc Thesis, University of Baghdad, Iraq.
- 16- **Rasheed** R O. (1994). A limnological study on some water systems in Erbil province. Iraq. M.Sc. Thesis. Univ. of Salahaddin-Erbil, Iraq.
- 17- **Al-Saadi**, H. A. ; Hadi, R. A. M. and Al-Lami, A. A. (1989). Limnological studies on some marsh areas in southern Iraq . Limnologia, 20 (2).
- 18- **Hashim**, N. N. (2010). Investigation of Cadmium and Mercury in Water,Sediments and Some Benthic Invertebrates at section of Tigris River in Baghdad City.Ms.C. Thesis, college of Science, Baghdad University: 125 PP.

- 19- Afzal,S;** Ahmad, I.;Younas, M.;Zahyd, M.; Khan,M.; Ijaz,A.andAli, K. (2000). Study of water quality of Hydiadrain , India. Pakistan. Environ-int. 29(1-2):87-96.
- 20- Siliem,** T. A. E. (1995): primary productivity of the Nile in barrage area. Menofiya J. Agric. Res.20 (4): 1687-1701.
- 20- Al-Helaly,** S. H. E. (2010). An investigation of some heavy metals in water, sediment and some biota of Al-Gharraf River, south of Iraq. M.Sc. Thesis. College of science, University of Baghdad.
- 21- Abed Al-Razzaq** H. T.(2011) Effect of Domestic Wastewater from Pumping Station of Al-Kadimiya on Ecological Properties of Tigris River. MSc. Thesis Collage of science, Baghdad university.
- 22- World Health Organization (WHO),**(2006). Guidelines for the safe use of wastewater, excreta and gray water: Wastewater use in agriculture. Volume II. France: 222pp.
- 23- Al- Nimma,** B. A. B. (1982). A study on the limnology of the Tigris and Euphrates rivers , M. Sci. Thesis Salahddin Univ. , Iraq.
- 24- Al-Saadi,** H. A.; Al-Lami, A. A.; Kassim, T. I. (1999) studies to ecological characteristics of upper Tigris and Euphrates Rivers and its correlation with fishery progress in Iraq. J. Environmental and sustainable progress researches, Vol.2, No.2- 1420. p. 24-31. (in Arabic)
- 25- Attee,** R. S. (2004) water characteristics in the Shatt Al-Arab river and the main drainage. Ph.D thesis, Basrah University. pp-128 (in Arabic)
- 26- USEPA,** (1999): National recommended water quality criteria - correction -United State Environmental Protection Agency EPA 822- Z-99-001;25p
- 27- Sawyer,** C.N. and McCarthy, P.L. (1978). Chemistry for environmental engineering. 3rd.Ed. McGraw-Hill Book Company. Singapore. 532pp.

- 28- Fahad, K. K.** (2005). Ecological survey for southern sector of Al-Garaf River, southern Iraq. Ph. D. Thesis. Collage of Agriculture, University of Basrah.
- 29- Sadek and Kamel** (2007). Seasonal variations in abiotic ecological conditions in Al-Garaf canal one of the main Tigris branches at Thi Qar province. J. Coll. Sci. Univ. Basrah :3-6
- 30- Lind, O. T.** (1979). Hand book of common methods in limnology. 2nd .Ed. London (109) pp .
- 31- Al-Gaffily, A. A. K.** (1992) Study of algae in Razzaza lake. MSc thesis. Baghdad University. p-102. (in Arabic)
- 32- Hassan, F. M.** (1997). A limnological study on Hilla river . Al- Mustainsiria J. Sci, 8 (1) : 22-30
- 33- AL-Saadi, H.A.; AL-Tamimi, A.N. and AL-Ghafily, A.A.** (1998) on the limnological features of Razzazah Lake, Iraq . Mutah J. for Research and Studies .3(2):111-127.
- 34- Al-Nimrawee, A.M.R.** (2005). The biodiversity of zooplankton and benthos invertebrates in Tigris and Euphrates River, Central Iraq. Ph. D. Thesis. Collage of Science, University of Baghdad.
- 35- Ajit M. Kalwale and Padmakar A. Savale** (2012). Determination of Physico-Chemical Parameters of Deoli Bhorus Dam water, Advances in Applied Science Research, 3 (1):273-279.
- 36- Al-Kuraishi, R.A.** (2011). A study of the effects of some ecological factors of Kut Dam on the Benthic Invertebrates of River Tigris. MSc. Thesis. Collage of science, Baghdad university.
- 37- Abed Al-Razzaq H. T.** (2011) Effect of Domestic Wastewater from Pumping Station of Al-Kadimiya on Ecological Properties of Tigris River. MSc. Thesis Collage of science, Baghdad university.
- 38- Joseph, N. C.; Eddy, A. O.; Elijah, J. P. and Ikechukwu, O. N. E.** (2011). Physicochemical evaluation of the effects of total suspended solids, total dissolved solids and total hardness concentrations on the water samples in Nsukka Town, Enugu State of Nigeria. Amr.J. Sci. 7 (5):827-836.
- 39- AL-Zamili, H.F.** (2007). Monthly variations of some environmental factors of AL-Gharraf River, J. of Dhi Qar, 3 (3) :17-2

- 40- Kathy, M.T.** (2008). Study of some physical and chemical properties of water and sediments of AL-Gharraf River, M.Sc. Thesis Collage of science, The-Qar University, 79p. (In Arabic).
- 41- Bilotta, G.S., Brazier, R.E.**, 2008. Understanding the influence of suspended solids on water quality and aquatic biota. *Water Res.* 42(3), 2849–2861.
- 42- Al-Saadi, H. A. ; Hadi, R. A. M. and Al-Lami, A. A.** (1989). Limnological studies on some marsh areas in southern Iraq . *Limnologica*, 20 (2).
- 43- Al-Fatlawy, Y.F.K.** (2007). Study the Drinking Water Quality of Some Baghdad Drinking Water Treatment Stations. Ph. D. Thesis. University of Baghdad. (In Arabic).
- 44- Ali, L. A.** (2007). A study of macroinvertebrates community in the middle sector of Greater Zab River, Iraq. Ph.D. Thesis. Univ. of Baghdad. Iraq.
- 45- Al-Lami, A.A. and Kassim, T. I. And Dulymi, A.A.** (1999). A limnological study on Tigris River. Iraq . *The Sci. J, IAEC.* 1:83.98.
- 46- Al-Ani, S. A.** (2002). Effect of Diyala River on the chemical and physical properties of Tigris River in southern Baghdad region. Ph.D. Thesis, college of Edu. Ibn- Al- Haithem, University of Baghdad: 170 PP.
- 47- Maulood, B. K. ; Al-Azzawi, M. N. and Saadalla, H. A.** (1994). An ecological study on the Tigris river pre and after crossing Baghdad. *J. Coll. Educ. Univ. Baghdad* 5 (1) : 43-50.
- 48- Al-Temymy, A.F.SH.** (2004). A microbiological and ecological study for the Tigris and Dyalah at the south of Iraq .MSc. Thesis, College of Science , Baghdad University. (In Arabic).

التغيرات الشهرية لبعض الصفات الفيزيائية والكيميائية لمياه نهر الغراف في مدينة الحي

وسام ثامر المياح و محمد نافع العزاوي

قسم علوم الحياة / كلية العلوم / جامعة بغداد

بغداد - العراق

الخلاصة:

درست التغيرات الشهرية في الخصائص الفيزيائية والكيميائية للمياه لتقييم نوعية مياه نهر الغراف بيئياً للمدة من تشرين الاول 2012 إلى نيسان 2013. حيث يقع نهر الغراف في الجزء الجنوبي الشرقي من العراق و تحيط فيه مساحات شاسعة وخصبة من الأراضي الزراعية. تم اختيار ثلاثة محطات للدراسة ، تقع المحطة الأولى على بعد 2 كم من مدينة الحي لتكون محطة السيطرة. وتقع المحطة الثانية على بعد 2 كم من المحطة الأولى حيث تمثل منطقة الدراسة .أما المحطة الثالثة فتقع على بعد 4 كم من المحطة الثانية بعد اجتياز النهر لمدينة الحي. تم أخذ العينات شهرياً وبواقع نموذجين لكل شهر.الغرض قياس قيم و تراكيز مؤشرات الماء وتراوحت التراكيز الشهرية كالآتي: درجة حرارة الهواء (16 الى 32) م⁰ و الماء (11 الى 26) م⁰ و الاس الهيدروجيني (7.05 الى 8.35) والاكسجين الذائب (7.04 الى 10.38) ملغم/ لتر والمتطلب البايولوجي للاوكسجين (0.90 الى 7.01) ملغم/ لتر و عكورة المياه 31.0 (الى 177.0) وحدة عكورة والمواد الصلبة الذائبة (550.78 الى 1108) ملغم/ لتر والمواد الصلبة العالقة (38 الى 636.5) ملغم/ لتر والتوصيلية الكهربائية (827 الى 1558) مايكروسيمنز والملوحة (0.54 الى 0.99) جزء بالألف والعسرة الكلية (275 الى 500) ملغم/ لتر والكلورايد (89.02 الى 184.6) ملغم/ لتر والنترات (5.7 الى 15.76) ملغم/ لتر والفوسفات (0.15 الى 0.5) ملغم/ لتر .