



Photoformation of reactive oxygen species and their potential to degrade highly toxic carbaryl and methomyl in river water

Aly Derbalah^{a, b}, Michael Sunday^{a, c}, Ryota Kato^a, Kazuhiko Takeda^a, Hiroshi Sakugawa^{a, *}

^a Graduate School of Biosphere Science, Hiroshima University, 1-7-1, Kagamiyama, Higashi-Hiroshima, Japan


^b Pesticides Chemistry and Toxicology Department, Faculty of Agriculture, Kafrelsheikh University, Kafrelsheikh, 33516, Egypt

^c Department of Chemistry, Federal University of Technology Akure, P.M.B 704, Ondo State, Nigeria

HIGHLIGHTS

- Reactive oxygen species ($^1\text{O}_2$ and $\cdot\text{OH}$) photogenerated in Kurose river water.
- Generation rate of $^1\text{O}_2$ was higher by two order than $\cdot\text{OH}$ in Kurose river water.
- The reaction rate constants of carbaryl and methomyl with $\cdot\text{OH}$ were several orders higher than $^1\text{O}_2$.
- Contribution of direct photolysis was very low compared to indirect photolysis in river water.
- $\cdot\text{OH}$ was the ROS largely responsible for the degradation of carbaryl and methomyl.

GRAPHICAL ABSTRACT



ARTICLE INFO

Article history:

Received 20 August 2019

Received in revised form

22 November 2019

Accepted 23 November 2019

Available online 25 November 2019

Handling Editor: Jun Huang

Keywords:

Hydroxyl radicals

Singlet oxygen

Methomyl

Carbaryl

Photolysis

Water

ABSTRACT

Reactive oxygen species (ROS) including singlet oxygen ($^1\text{O}_2$) and hydroxylradicals ($\cdot\text{OH}$) photogenerated in natural waters play important roles in indirect photolysis of man-made pollutants. This study was conducted to investigate how the generation of these two ROS influences the degradation of two highly toxic insecticides (methomyl and carbaryl) in river water. To accomplish this, the reaction rate constants of $^1\text{O}_2$ and $\cdot\text{OH}$ with carbaryl and methomyl were determined; the degradation rate constants of the tested insecticides in ultrapure water (direct photolysis) and in river water in the presence and absence of $^1\text{O}_2$ and $\cdot\text{OH}$ scavengers were also measured. The rate constants for the reaction of $\cdot\text{OH}$ with carbaryl and methomyl were found to be $(14.8 \pm 0.64) \times 10^9$ and $(4.68 \pm 0.52) \times 10^9 \text{ M}^{-1} \text{ s}^{-1}$, respectively. The reaction rate constant of $^1\text{O}_2$ with carbaryl $(2.98 \pm 0.10) \times 10^5 \text{ M}^{-1} \text{ s}^{-1}$, was much higher than that of methomyl ($< 10^4 \text{ M}^{-1} \text{ s}^{-1}$). Indirect photolysis by $\cdot\text{OH}$ accounted for 63% and 62%, while $^1\text{O}_2$ accounted for 26% and 30% and direct photolysis accounted for 1.4% and 7% of methomyl and carbaryl degradation, respectively. The high degradation rate in river water demonstrated by both insecticides suggests that indirect photolysis mediated by $\cdot\text{OH}$ is an important means of their degradation in river water. In addition, kinetic calculations of $\cdot\text{OH}$ -mediated degradation rate constants of the compounds agrees with their experimentally-determined values thereby confirming the importance of $\cdot\text{OH}$ towards their degradation.

© 2019 Elsevier Ltd. All rights reserved.

* Corresponding author.

E-mail address: hsakuga@hiroshima-u.ac.jp (H. Sakugawa).

Credit author statement

Aly Derbalah: Doing experiments, Writing, Publishing, Analysis
Michael Sunday: Doing experiments, Writing, Reviewing
Ryota Kato: Doing experiments, **Kazuhiko Takeda:** Sampling Supervision
Hiroshi Sakugawa: Reviewing, Supervision.

1. Introduction

Pesticides have been used intensively worldwide to protect crops from pests and diseases in order to maintain adequate productivity so that food is more affordable. Pesticides will continue to be used as an effective means of controlling pests and increasing agricultural production (Alhousari, 2011). Methomyl and carbaryl (Fig. 1S) are a carbamate compounds widely used to control several insects in many different types of crops (Tomlin, 2003; EPA, 2004) and considered very toxic to humans, aquatic organisms, and other beneficial organisms such as bees and birds (Street, 1981; Barceló et al., 1996; McDuffie, 2001; Drea et al., 2012; Bazrafshan et al., 2017). Due to their extensive use, methomyl and carbaryl have been detected in surface and ground water at various concentrations (Chowdhury et al., 2012; Chattoraj et al., 2014; Struger et al., 2016).

In the aquatic environment, pesticides are transformed by photo- and biodegradation (Derbalah et al., 2013, 2014; 2016; Katagi, 2018). However, photodegradation (direct and indirect) is one of the most important abiotic transformations for pesticides in the aquatic environment (Katagi, 2018). Several reactive species, including hydroxyl radicals ($\cdot\text{OH}$), singlet oxygen ($^1\text{O}_2$), triplet oxygen, chromophoric organic matter in the triplet state, superoxide ions and hydroperoxyl radicals ($\text{O}_2\cdot^-/\text{HO}_2\cdot$), and carbonate radicals ($\cdot\text{CO}_3^-$), can be produced upon sunlight irradiation of components such as dissolved organic matter (DOM), nitrate, nitrite, and Fe(III) present in natural waters (Zepp et al., 1981; Draper and Crosby, 1984; Haag and Hoigné, 1986; Richard and Canonica, 2005; Vione et al., 2009). These species play important roles in the chemical breakdown of organic pollutants such as pesticides in water (Street, 1981; Haag and Hoigné, 1986; McDuffie, 2001; Vermilyea and Voelker, 2009; Arakaki et al., 2010).

Among these species, $\cdot\text{OH}$ and $^1\text{O}_2$ contribute significantly to the indirect photodegradation of a wide range of organic pollutants. $\cdot\text{OH}$ is perhaps the most reactive oxygen specie, exhibiting high reaction rate constants with a wide range of organic and inorganic pollutants in natural waters (Buxton et al., 1988). Its high reaction rate constants of approximately $10^9 \text{ M}^{-1} \text{ s}^{-1}$ (Vermilyea and Voelker, 2009) with organic pollutants enables it to contribute to the breakdown of organic pollutants such as pesticides that are considered relatively stable and cannot be destroyed by biological processes or direct photolysis (Mabury and Crosby, 1996; Armbrust, 2000; Alhousari et al., 2010; Dell'Arciprete et al., 2010; Arakaki et al., 2010).

$^1\text{O}_2$ is formed by energy transfer from the triplet excited states of DOM to dissolved molecular oxygen (Zepp et al., 1981; Haag and Hoigné, 1986; Richard and Canonica, 2005). The reactivity of $^1\text{O}_2$ with organic compounds, including pesticides, is relatively lower compared to the reactivity of $\cdot\text{OH}$. Nevertheless, its higher steady-state concentration in the aquatic environment, which can be three to four orders of magnitude greater than that of $\cdot\text{OH}$, gives it a competitive advantage to participate in pesticides degradation. Therefore, this may also represent an important degradation pathway for pesticides.

An understanding of the fates of pesticides is essential for rational decision-making regarding their authorization.

Furthermore, information regarding the possible pathways and scenarios for pesticide degradation in aquatic environments is crucial; ensuring that the period for which they persist in the environment and the potential risks they pose to humans and aquatic organisms can be safely predicted (Alhousari, 2011). Due to the high toxicity and frequent detection of both carbaryl and methomyl in water (Street, 1981; Barceló et al., 1996; McDuffie, 2001; Wilsont and Foos, 2006; Chowdhury et al., 2012; Chattoraj et al., 2014; Struger et al., 2016), there is a need to investigate their photodegradation in river water and also unravel the contributions of direct photolysis and/or indirect photolysis (mediated by ROS) under environmentally-relevant irradiation conditions. To the best of our knowledge, such investigation has not been conducted before now. Hence, this study was conducted to evaluate the photodegradation of methomyl and carbaryl, as well as to obtain their half-lives under environmentally relevant irradiation conditions in river water. The contributions of $\cdot\text{OH}$ and $^1\text{O}_2$ to the degradation of the two compounds in river water were also determined.

2. Materials and methods

2.1. Reagents and chemicals

Acetonitrile and benzene were purchased from Nacalai Tesque, Kyoto, Japan (HPLC grade > 99.5%). Phenol methomyl (99.9%), carbaryl (99.9%), and potassium hydrogen phthalate were obtained from Sigma-Aldrich (Japan). Sodium nitrite, sodium nitrate, sodium chloride, sodium sulfate, iron (II) sulfate, sulfanilamide, rose bengal (RB), and furfuryl alcohol (FFA) were purchased from Katayama Chemical Industries (guaranteed grade) (Japan). Hydrogen peroxide (ca. 30%) was obtained from Wako Pure Chemical Industries (Japan). The 2-nitrobenzaldehyde (2-NB) and *N*-1-naphthyl ethylenediamine dichloride were purchased from Tokyo Kasei Kogyo (Japan). Disodium terephthalate (TP) and 2-hydroxyterephthalic acid (HTP) were obtained from Tokyo Chemical Industry Co. Ltd. (Japan). Stock and working solutions of the examined insecticides were prepared in ultrapure water only, without solvents.

2.2. Water samples

Surface river water samples were collected in June 2018 from Izumi Station on the Kurose River (chemical composition shown in Table 1S in Higashi-Hiroshima, Japan. Samples were collected into clean glass 1-L capacity bottles, then filtered through fiberglass filters (Advantech, Tokyo, Japan, 0.45 μm nominal rating) for further treatment. Samples were stored in brown borosilicate bottles at 4 °C to prevent biological degradation until analysis (within a week).

2.3. ROS determination

Determination of $\cdot\text{OH}$ and $^1\text{O}_2$ generated during irradiation of water samples was done using chemical probes. For these determinations, irradiation was done using a solar simulator consisting of a lamp housing (WACOM Co., Saitama, Japan HX-500) equipped with a 500 W Xe lamp (WACOM Co., KXL-500F) and an optical filter (HOYA Co., Tokyo, Japan ultraviolet (UV)-31) with a transmission wavelength limit of 310 nm. A detailed description of the solar simulator has been reported elsewhere (Nakatani et al., 2007). The photoformation rates and steady-state concentrations of $\cdot\text{OH}$ and $^1\text{O}_2$ were normalized to 2-NB degradation rate of


0.0093 s⁻¹ as previously described (Takeda et al., 2014).

The amount of photoformed ·OH in the river water sample was determined using benzene as a chemical probe (Takeda et al., 2004). Benzene reacts with ·OH at a high bimolecular reaction rate constant of 7.8 × 10⁹ M⁻¹ s⁻¹ to produce phenol (Buxton et al., 1988). A sample containing 1.2 mM benzene was irradiated for 40 min while aliquots were withdrawn at 10 min intervals and analyzed for phenol using high-performance liquid chromatography (HPLC). The photoformation rate, scavenging rate constant, and steady-state concentration of ·OH were determined and are detailed in the supplementary information (SI), section S1.

¹O₂ photogenerated in the irradiated samples was measured using FFA as a probe. It reacts with ¹O₂ and degrades following first-order degradation kinetics, (Haag and Hoigné, 1986). Samples were spiked with FFA (final concentration, 100 μM) and irradiated under the solar simulator, during this time aliquots of the reaction mixture were obtained and analyzed for FFA using HPLC. The detailed procedure is available under S1. The ¹O₂ steady-state concentration ([¹O₂]_{ss}) was determined by dividing the observed first-order degradation rate constant (k) of FFA in the irradiated sample by the reaction rate constant of FFA with ¹O₂ (k_{FFA,O₂}¹), which was given as 1.09 × 10⁸ M⁻¹ s⁻¹ (Haag et al., 1984). By considering water as the main scavenger of ¹O₂ in natural waters, with a k_d of 2.5 × 10⁵ s⁻¹ (Rodgers and Snowden, 1982), the photoformation rate of ¹O₂ (R¹O₂) was obtained as a product of the [¹O₂]_{ss} and k_d.

2.4. Reaction rate constant of target insecticides with ·OH

The reaction rate constants between the tested insecticides and ·OH were determined based on competition kinetics with TP as the ·OH probe (Yang and Guo, 2001; Louit et al., 2009; Charbouillot et al., 2011; Bekdeser et al., 2012; Takeda et al., 2017a,b). The ·OH was generated using a high-power 365 nm UV-light-emitting diode (LED) (NS365L-6SMG, Nitride Semiconductors, Naruto, Japan) and 0.1 μM nitrite aqueous solution (Takeda et al., 2018), and was detected with 0.05 mM TP as shown in equation (1).



In the irradiated sample solution containing TP, nitrite, and insecticide, the generated ·OH competitively reacted with the TP and insecticides. The TP reacted with ·OH quantitatively to generate a strongly fluorescent product, HTP (Yang and Guo, 2001; Louit et al., 2009; Charbouillot et al., 2011; Bekdeser et al., 2012; Takeda et al., 2017a). The reaction rate constant was calculated by equation (2):

$$\frac{F_0}{F} = 1 + \frac{k_{X,\text{OH}}[C]}{k_{\text{TP},\text{OH}}[\text{TP}]} \quad (2)$$


where $k_{\text{TP},\text{OH}}$ and [TP] are the reaction rate constant and concentration of TP, respectively, and $k_{X,\text{OH}}$ and [C] are the reaction rate constant and concentration of the insecticide, respectively. Additionally, F is the fluorescence signal of HTP in the presence of the insecticide while F₀ is the fluorescence signal of HTP without insecticide, i.e., [C] = 0. Equation (2) indicates that a plot of F₀/F against [C] at constant [TP] should give a straight line with a y-intercept of 1 and a slope of $k_{X,\text{OH}}/(k_{\text{TP},\text{OH}} [\text{TP}])$. The reaction rate constant of the insecticides with ·OH ($k_{X,\text{OH}}$) can then be simply calculated from the slope of this plot and the values of $k_{\text{TP},\text{OH}}$ and [TP]. We selected a $k_{\text{TP},\text{OH}}$ of 4.0 × 10⁹ M⁻¹ s⁻¹ (Charbouillot et al., 2011) to calculate the rate constants of the target insecticides.

To determine the reaction rate constants, a quartz photochemical reaction cell (20 mm in diameter and 12 mm thick (optical path)) was used. The UV-LED was placed 10 mm above the top

surface of the reaction cell, after which the sample solution was irradiated for 10 min. The HTP formed was analyzed using HPLC as described by Takeda et al. (2018).

2.5. Reaction rate constant of target insecticides with ¹O₂

The reaction rate constants of ¹O₂ with carbaryl and methomyl were determined by monitoring the degradation rates of methomyl and carbaryl as they react with photogenerated ¹O₂, using RB as a sensitizer (equations (3) and (4)) (Vione et al., 2011; Ruggeri et al., 2013). The reaction rates of the insecticides with ¹O₂ were then determined as shown in equation (5):



$$\text{Rate of target insecticides degradation (Rdeg)} = k_{\text{insecticides},1\text{O}_2} [\text{insecticides}] [{}^1\text{O}_2] \quad (5)$$

where $k_{\text{insecticides},1\text{O}_2}$ is the reaction rate constant of insecticides with ¹O₂. At constant [¹O₂], R_{deg} is directly proportional to the insecticide concentration. Therefore, as the insecticide concentration increases, R_{deg} is expected to increase. Moreover, when insecticides react with ¹O₂, R_{deg} can also be expressed as the product of the rate of ¹O₂ generation (R¹O₂) and the fraction (F) of ¹O₂ that reacts with the insecticides. This is expressed in equations (6)–(9) below.

$$R_{\text{deg}} = R_{{}^1\text{O}_2} \times (F) \quad (6)$$

$$F = \frac{k_{\text{insecticides},1\text{O}_2} [\text{insecticides}]}{k_d + k_{\text{insecticides},1\text{O}_2} [\text{insecticides}]} \quad (7)$$

where k_d is the dissociation rate constant of ¹O₂ in water = 2.5 × 10⁵ s⁻¹ as reported previously (Rodgers and Snowden, 1982).

$$R_{\text{deg}} = R_{{}^1\text{O}_2} \times \frac{k_{\text{insecticides},1\text{O}_2} [\text{insecticides}]}{k_d + k_{\text{insecticides},1\text{O}_2} [\text{insecticides}]} \quad (8)$$

At low concentrations of insecticides (μM range), $k_{\text{insecticides},1\text{O}_2} [\text{insecticides}] \ll k_d$, giving rise to equation (9):

$$R_{\text{deg}} = R_{{}^1\text{O}_2} k_d^{-1} k_{\text{insecticides},1\text{O}_2} [\text{insecticides}] \quad (9)$$

From equation (9), a plot of R_{deg} against [insecticides] at a constant ¹O₂ generation (R¹O₂) can be generated that gives a slope = $R_{{}^1\text{O}_2} k_d^{-1} k_{\text{insecticides},1\text{O}_2}$. Because k_d is known, R¹O₂ can be obtained using FFA under the same experimental conditions.

R¹O₂ was determined from the initial degradation rate of FFA (R_{FFA}) as shown in equation (10) below.

$$R_{\text{FFA}} = R_{{}^1\text{O}_2} \times \frac{k_{\text{FFA},1\text{O}_2} [\text{FFA}]}{k_d + k_{\text{FFA},1\text{O}_2} [\text{FFA}]} \quad (10)$$

where $k_{\text{FFA},1\text{O}_2}$ is the reaction rate constant between FFA and ¹O₂, which is 1.09 × 10⁸ M⁻¹ s⁻¹ (Haag and Hoigné, 1986). A detailed experimental procedure for the determination of R_{deg} and R¹O₂ is provided in SI S2.

2.6. Photodegradation of carbaryl and methomyl in ultrapure water and river water

Carbaryl and methomyl solutions (1 mg/L) in ultrapure water were irradiated under the solar simulator using a Pyrex filter to investigate the contributions of direct photolysis to their degradations at a buffered pH value of 6.8. Dark controls without light were performed to account for the degradation under dark conditions. The degradation under dark conditions was negligible and the data are not shown. Photochemical experiments for direct degradation rates were conducted with a solar simulator as described by Takeda et al. (2014).

Photodegradation studies of carbaryl and methomyl (1 mg/L) in river water in the absence and presence of 1% methanol ($\cdot\text{OH}$ scavenger) (Motohashi and Saito, 1993) and 1 mM NaN_3 ($^1\text{O}_2$ scavenger) (Miskoski and García, 1993) were conducted using a solar simulator that consisted of a lamp housing (WACOM Co., HX-500) equipped with a 500 W Xe lamp (WACOM Co., KXL-500F) and an optical filter (HOYA Co., UV-31) with a transmission wavelength limit of 310 nm (Dell'Arciprete et al., 2010). Samples were withdrawn at intervals and analyzed by HPLC (described above) to monitor the degradation of the insecticides. The degradation rate constant (k) was obtained as the slope of the plot of the natural logarithm of concentration ($\ln C_t/C_0$) against irradiation time, where C_t was the concentration of the compound at time (t) and C_0 was the initial concentration of the compound. The obtained k values were normalized to a 2-NB degradation rate of 0.0093 s^{-1} as previously described (Arakaki et al., 1999). This experiment was conducted in triplicate. The contribution of each ROS ($\cdot\text{OH}$ and $^1\text{O}_2$) in the degradation of the tested insecticides in water was estimated by equation (11):

$$\text{ROS contribution} = (\text{J}_{\text{river}} - \text{J}_{\text{scavenger}}) / \text{J}_{\text{river}} \times 100 \quad (11)$$

where ROS contribution represents the contribution (%) of $\cdot\text{OH}$ and $^1\text{O}_2$ in the degradation of each insecticide, J_{river} is the degradation rate of each insecticide in river water and $\text{J}_{\text{scavenger}}$ is the degradation rate of each insecticide in the presence of $\cdot\text{OH}$ or $^1\text{O}_2$ scavengers. The contribution (%) of direct photolysis in the degradation of each insecticide was estimated by equation (12):

$$\text{Direct photolysis} = (\text{J}_{\text{water}} / \text{J}_{\text{river}}) \times 100 \quad (12)$$

where J_{water} is the degradation rate of each insecticide in ultrapure water.

2.7. Calculation of photochemical half-lifetimes

The photochemical half-lives for the tested insecticides as mediated by each ROS were calculated using equation (13) (Nakatani, 2004):

$$t_{1/2} = \frac{\ln 2}{F_{\text{solar}} \times (k_{\text{direct}} + k_{\text{insecticide,ROS}} [\text{ROS}]_{\text{ss}})} \quad (13)$$

where k_{direct} is the direct photolysis rate constant in Milli-Q water, $k_{\text{insecticide,ROS}}$ is the reaction rate constant of each insecticide with ROS ($\cdot\text{OH}$ or $^1\text{O}_2$), and $[\text{ROS}]_{\text{ss}}$ is the steady-state concentration of the ROS. Additionally, F_{solar} is a light intensity factor (taken as 0.1) to account for the diurnal and seasonal changes in the solar radiation intensity. F_{solar} was obtained as a product of 0.3 (a factor used to indicate 8 h of daily sunlight exposure) and 1/3 (based on winter sunlight intensity being approximately one-third that of summer according to the solar actinic flux at the Earth's surface) (Trapp and Matthies, 1995; Diamond et al., 2001; Takeda et al., 2014; Kaonga

et al., 2016).

2.8. Data analysis

For analysis of variance (ANOVA) of obtained data, XLSTAT PRO statistical analysis software (Addinsoft) was used. Fisher's least significant difference (LSD) test was used to separate the mean of each treatment. All analyses were performed at a significance value of $P \leq 0.05$.

3. Results and discussion

3.1. $\cdot\text{OH}$ and $^1\text{O}_2$ generation in river water

The generation of $\cdot\text{OH}$ and $^1\text{O}_2$ in river water was investigated using benzene and FFA, respectively, as selective probes. As shown in Table 1, the photoformation rate ($R^1\text{O}_2$) and steady-state concentration $[^1\text{O}_2]_{\text{ss}}$ in the river water were $7.9 \times 10^{-8} \text{ M s}^{-1}$ and $3.1 \times 10^{-13} \text{ M}$, respectively. These values were two orders higher than those of the $\cdot\text{OH}$ photoformation rate (R_{OH}) and steady-state concentration $[\cdot\text{OH}]_{\text{ss}}$ of $1.76 \times 10^{-10} \text{ M s}^{-1}$ and $1.2 \times 10^{-15} \text{ M}$, respectively. The reported values were normalized against natural sunlight intensity using 2-NB as a chemical actinometer.

In this study, the formation rate of the $\cdot\text{OH}$ radicals in Kurose River water was on the same order as that reported by Takeda et al. (2004). The photoproduced $\cdot\text{OH}$ in water from the Kurose River was large due to the concentrations of nitrate and nitrite (Takeda et al., 2004). The $[^1\text{O}_2]_{\text{ss}}$ values obtained in this study agree with the typical values (10^{-13} M) reported in river water samples (Peterson et al., 2012). The steady-state concentration of ROS is a ratio of the photoformation rate and the scavenging rate constant (SRC). The SRC of $\cdot\text{OH}$ in this study ($1.47 \times 10^5 \text{ s}^{-1}$) is on the same order as the k_d of $^1\text{O}_2$ in water ($2.5 \times 10^5 \text{ s}^{-1}$) (Rodgers and Snowden, 1982). Therefore, the higher steady-state concentration of $^1\text{O}_2$ than that of $\cdot\text{OH}$ in river water is because of the higher photoproduction rate of $^1\text{O}_2$. Moreover, the high reactivity and short lifetime of $\cdot\text{OH}$ cause its steady-state concentrations in natural waters to be generally low (Takeda et al., 2004). In addition, $\cdot\text{OH}$ is efficiently consumed by natural DOM, which limits its steady-state concentration in water illuminated by sunlight (Richard and Canonica, 2005). Unlike $\cdot\text{OH}$, which has more than one source (NO_3^- , NO_2^- , DOM, H_2O_2) and sink (DOM, Br^- , Cl^-) (Takeda et al., 2004), the main source of $^1\text{O}_2$ in natural waters is chromophoric DOM (Peterson et al., 2012), while its primary sink is deactivation by water molecules.

3.2. Reaction rate constants of the tested insecticides with $\cdot\text{OH}$ and $^1\text{O}_2$

To examine the capacity of $\cdot\text{OH}$ and $^1\text{O}_2$ to degrade the examined insecticides, the reaction rate constants of methomyl and carbaryl with the two ROS were measured. The reaction rate constants of $\cdot\text{OH}$ with the insecticides were determined by the competition kinetics method using TP as a reference compound as described by Takeda et al. (2017a). As shown in Table 2, the reaction rate constant of $\cdot\text{OH}$ with carbaryl ($14.8 \times 10^9 \text{ M}^{-1} \text{ s}^{-1}$) was approximately three times higher than that of methomyl ($4.68 \times 10^9 \text{ M}^{-1} \text{ s}^{-1}$). The electron-rich aromatic rings present in carbaryl would be the most likely primary site for addition reactions with hydroxyl radicals. Hydrogen abstraction could also contribute to the reactivity (Scully and Hoigné, 1987). According to equation (2), a plot of F_0/F against insecticide concentrations $[X]$ with an intercept of one is required to determine the reaction rate constant. Fig. 2S shows a plot of F_0/F against the insecticide concentrations $[X]$ that were used to determine the reaction rate constants of the insecticides with $\cdot\text{OH}$. An intercept of

Table 1
Steady-state concentration and photoformation rate of ROS in water from the Kurose River.

Reactive oxygen species	Steady-state concentration (M)	Photoformation rate (Ms ⁻¹)	Scavenging rate (s ⁻¹)
·OH	(1.2 ± 0.01 ^a) × 10 ⁻¹⁵	(1.76 ± 0.01 ^a) × 10 ⁻¹⁰	(1.47 ± 0.01 ^a) × 10 ⁵
¹ O ₂	(3.1 ± 0.02 ^b) × 10 ⁻¹³	(7.9 ± 0.03 ^b) × 10 ⁻⁸	(*2.5 ± 0.02 ^b) × 10 ⁵

* Deactivation rate constant of ¹O₂ by water as reported by Rodgers and Snowden (1982).

*Statistical comparisons were made among treatments within a single column.

*The different letters represent significant differences using Fisher's LSD test at P ≤ 0.05.

*Each mean value came from three replicates.

Table 2
Reaction rate constants of the tested insecticides with ·OH and ¹O₂.

ROS	Reaction rate constants (M ⁻¹ s ⁻¹)	
	Methomyl	Carbaryl
·OH	(4.68 ± 0.25) × 10 ⁹	(14.8 ± 0.12 ^a) × 10 ⁹
¹ O ₂	< 10 ⁴	(2.98 ± 0.10 ^b) × 10 ⁵

*Statistical comparisons were made among treatments within a single column.

*The different letters represent significant differences using Fisher's LSD test at P ≤ 0.05.

*Each mean value came from three replicates.

approximately one as expected from equation (2) was obtained.

The reaction rate constants of ¹O₂ with the insecticides were determined by monitoring the degradation of the insecticides at varying concentrations in the presence of high [¹O₂]_{ss} generated using RB as a sensitizer (Vione et al., 2011; Ruggeri et al., 2013). A plot of the R_{deg} of carbaryl as a function of its concentration is shown in Fig. 3S. According to equation (9), a plot of R_{deg} against [insecticide] gives a slope = R_{1O2} k_d⁻¹ k_{insecticides} ¹O₂. The slope obtained for the carbaryl detection experiment (Fig. 3S) was 2.11 × 10⁻⁵ M⁻¹ s⁻¹. The R_{1O2} obtained using FFA in the experiment instead of carbaryl was 1.78 × 10⁻⁵ M⁻¹ s⁻¹, while the k_d was taken to be 2.5 × 10⁵ s⁻¹ (Rodgers and Snowden, 1982). Based on these values, the reaction rate constant of ¹O₂ with carbaryl was determined to be 2.98 × 10⁵ M⁻¹ s⁻¹. However, the reaction rate constant of ¹O₂ with methomyl was very low (<10⁴ M⁻¹ s⁻¹). This was because methomyl at varying concentrations did not undergo any significant degradation in the presence of high concentrations of ¹O₂. Analysis of methomyl did not reveal enough degradation to enable calculation of its reaction rate constant with ¹O₂. This suggests that the methomyl rate constant with ¹O₂ is much lower than that of ¹O₂ with carbaryl. ¹O₂ is reactive towards electron-rich compounds (Scully and Hoigné, 1987), and the absence of such electron-rich functional groups in methomyl explains why it showed a very minimal reaction with ¹O₂. The aromatic rings present in carbaryl may be responsible for its better reactivity with ¹O₂ compared with methomyl.

Our results showed that the reaction rate constant of carbaryl with ¹O₂ was approximately five orders of magnitude lower than its rate constant with ·OH. Similarly, the reaction rate constant of methomyl with ¹O₂ (which is much lower than that of carbaryl with ¹O₂) will also be several orders lower than its rate constant with ·OH. This great difference in the rate constants of the insecticides with the two ROS suggests the importance of ·OH in the photodegradation of the two compounds in the environment. The steady-state concentration of ¹O₂ in river water is usually higher (two orders in this study) than that of ·OH. Nevertheless, the rate constants of the insecticides with ·OH are far higher than those with ¹O₂. These higher rate constants are enough to offset the potential contribution associated with the higher steady-state concentration of ¹O₂.

3.3. Photodegradation of methomyl and carbaryl in ultrapure and river water

The direct photolysis of the tested insecticides (methomyl and carbaryl) in ultrapure water normalized to the intensity of natural sunlight is shown in Table 3. The data showed that the degradation rate of carbaryl (4.62 × 10⁻⁶ s⁻¹) was much higher than that of methomyl (6.27 × 10⁻⁷ s⁻¹). This suggests that methomyl is more stable than carbaryl when subjected to direct photolysis. Photolytic transformations of the tested insecticides are caused by the absorption of energy (photons) from light. By absorbing light energy, these pesticides are transformed through a number of chemical reactions including cleavage of chemical bonds, oxidation, and hydrolysis (Zepp and Cline, 1977; Clark, 1994; Larson et al., 1997).

The degradation rates of the target insecticides in river water, 6.5 × 10⁻⁵ s⁻¹ (for carbaryl) and 4.47 × 10⁻⁵ s⁻¹ (for methomyl), were generally higher than that of the degradation rate in ultrapure water (direct photolysis). This suggests the significant contribution of indirect photolysis mediated by ROS in the degradation of the target insecticides in river water. Hence, the contribution of ·OH and ¹O₂ to the photolysis of the target insecticides in river water was investigated by scavenger experiments. As shown in Fig. 1 and Table 3, the degradation rate of carbaryl in the absence of scavengers (6.5 × 10⁻⁵ s⁻¹) was higher than in the presence of 1% methanol (2.03 × 10⁻⁵ s⁻¹) and 1 mM NaN₃ (3.66 × 10⁻⁵ s⁻¹) as ·OH and ¹O₂ scavengers, respectively. Similarly, the degradation rate of methomyl in the absence of scavengers (4.47 × 10⁻⁵ s⁻¹) was higher than in the presence scavengers of ·OH (1.62 × 10⁻⁵ s⁻¹) and ¹O₂ (3.25 × 10⁻⁵ s⁻¹) as shown in Table 3 and Fig. 2. As shown in Fig. 3, photodegradation studies in the presence of scavengers revealed that ·OH accounted for 63 and 62% of the degradation of methomyl and carbaryl, respectively, while ¹O₂ accounted for 26 and 30% and direct photolysis accounted for 1.4 and 7.1%, respectively. The residual degradation percentage of methomyl (9.3%) and carbaryl (1%) may be because of the generation of other photosensitizers such as chloride and sulfate radicals (Lian et al., 2017). These findings demonstrate that indirect photolysis mediated by ·OH is an important means of degradation of the two insecticides in river water. Miller and Chin (2002) also reported that NOM and nitrate in conjunction acted as photosensitizers that resulted in the indirect photolytic degradation of carbaryl in wetland surface waters. Additionally, they suspected that the principal pathway occurs through the generation of ROS, which in turn are capable of reacting with the target pesticide.

3.4. Validation of ROS contributions from kinetic calculations

The involvement of the two ROS in the degradation of the insecticides was further investigated by comparing the experimentally-determined degradation rate constants of the insecticide, as mediated by each ROS, with the calculated degradation rate constants. The experimental values were obtained by deducting the degradation rate constants in the presence of a

Table 3
Degradation rate constants of the tested insecticides in different water matrices.

Water type	Degradation rate constant, k (s^{-1})	
	Methomyl	Carbaryl
Ultrapure water (direct photolysis)	$(6.27 \pm 0.31^a) \times 10^{-7}$	$(4.62 \pm 0.22^a) \times 10^{-6}$
River water (direct + indirect photolysis)	$(4.47 \pm 0.27^b) \times 10^{-5}$	$(6.50 \pm 0.34^b) \times 10^{-5}$
River water + NaN_3 ($^1\text{O}_2$ scavenger)	$(3.25 \pm 0.17^c) \times 10^{-5}$	$(4.10 \pm 0.15^b) \times 10^{-5}$
River water + methanol ($\cdot\text{OH}$ scavenger)	$(1.62 \pm 0.10^d) \times 10^{-5}$	$(2.03 \pm 0.10^c) \times 10^{-5}$

*Statistical comparisons were made among treatments within a single column.

*The different letters represent significant differences using Fisher's LSD test at $P \leq 0.05$.

*Each mean value came from three replicates.




Fig. 1. Degradation rate constants of carbaryl in river water in the presence and absence of $^1\text{O}_2$ (NaN_3) and $\cdot\text{OH}$ (methanol) as scavengers.




Fig. 2. Degradation rate constants of methomyl in river water in the presence and absence of $^1\text{O}_2$ (NaN_3) and $\cdot\text{OH}$ (methanol) as scavengers.

scavenger from the degradation rate constants obtained in the absence of the scavenger during irradiation of insecticide-spiked river water sample. The degradation rates due to direct photolysis were also deducted. The calculated degradation rate constants were obtained as a product of $k_{\text{insecticide, ROS}}$ and $[\text{ROS}]_{\text{ss}}$ for corresponding ROS - insecticide pair. Details have been provided in supplementary information (S3). The results (Table 2S) showed that, for $\cdot\text{OH}$ -mediated degradation of carbaryl, there was an agreement between the experimental degradation rate constants ($4 \times 10^{-5} \text{ s}^{-1}$) and the calculated rate constants ($1.8 \times 10^{-5} \text{ s}^{-1}$). Similarly, there was a considerable agreement between the experimental ($2.0 \times 10^{-5} \text{ s}^{-1}$) and calculated ($0.56 \times 10^{-5} \text{ s}^{-1}$) degradation rate constants for $\cdot\text{OH}$ -mediated degradation of methomyl. However, for $^1\text{O}_2$ -mediated degradation of the insecticides, agreements between the calculated and the experimentally-determined degradation rate constants could not




Fig. 3. Contribution of each ROS ($^1\text{O}_2$ and $\cdot\text{OH}$) and the direct photolysis to degradation of the tested insecticides in water. *Other oxidants (i.e., chloride and sulfate radicals, Lian et al., 2017).

be established (Table 2S). This disagreement could be due to the microheterogeneous distribution of $^1\text{O}_2$ between the DOM phase and bulk aqueous phase. It has been reported that $^1\text{O}_2$ in the DOM microenvironment could be several orders (up to three orders) of magnitude larger than in the aqueous phase (Latch and McNeil, 2006; Grandbois et al., 2008). The hydrophobic nature of carbaryl, as evidenced by its very low water solubility (4 mg/100 mL), suggests it may bind to DOM in river water. The binding of carbaryl to DOM may expose it to $[\text{O}_2]$ in the DOM microenvironment which is several orders of magnitude higher than the $[\text{O}_2]$ in the bulk aqueous environment. Under such circumstances, the $^1\text{O}_2$ -mediated degradation of the insecticide may proceed largely in the DOM region. Hence, the experimentally-determined degradation rate constants will differ significantly from the calculated values which are based on aqueous phase $[\text{O}_2]$. Regardless of the discrepancy between the calculated and experimental $^1\text{O}_2$ degradation rate constants, the agreement between the experimental and calculated degradation rate constants for $\cdot\text{OH}$ -mediated degradation of these insecticides confirms the role of $\cdot\text{OH}$ as the primary ROS responsible for their degradation in river water.

3.5. Half-lives in river water

The half-lives of the insecticides as mediated by each ROS ($\cdot\text{OH}$ and $^1\text{O}_2$) in river water were determined using equation (12). The determined half-lives were normalized to an irradiation condition (F_{solar}) of approximately 8 h (0.3 d) of daily sunlight and a 1/3 variation in yearly light intensity based on light intensity in winter

being 1/3 that of summer. As shown in Table 4, the calculated half-lives of carbaryl (3.6 d) in river water with $\cdot\text{OH}$ were shorter than that of the half-life with $^1\text{O}_2$ (17.06 d). Furthermore, the calculated half-life of methomyl in river water with $\cdot\text{OH}$ (13.3 d) was much lower than that with $^1\text{O}_2$ (>186 d). The shorter photochemical half-life of carbaryl could be attributed to its higher reactivity with $\cdot\text{OH}$ and $^1\text{O}_2$ and higher direct photolysis compared with methomyl. This higher reactivity of carbaryl may be because of its aromatic structure compared with the aliphatic structure of methomyl, which makes $\cdot\text{OH}$ more reactive with electron-rich aromatic organic compounds because of its electrophilicity and non-selective nature towards C–H bonds (Zepp et al., 1985; Buxton et al., 1988; Mabury and Crosby, 1996).

Overall, the results indicate that $\cdot\text{OH}$ specie play an important and major role in the breakdown of pollutants in water, including the insecticides investigated in this study, because of their high reaction rate constants toward a very wide range of organic and inorganic compounds (Bazrafshan et al., 2017). Because of their high reactivity with organic compounds, $\cdot\text{OH}$ species provide effective solutions to the breakdown of organic compounds with high stability and resistance to chemical, biological, and direct photochemical degradation (Walse et al., 2004; Richard and Canonica, 2005; Takeda et al., 2017a).

Finally, inspite of Kurose river water flow into Seto Inland sea, it is very difficult to discuss how much the methomyl and carbaryl are photodegraded when these insecticides in river water flow into seawater using the calculated half-lives. This is because the [ROS] in River water is completely different from [ROS] in seawater. In addition, the sample matrix of seawater is completely different from that of river water. Therefore, it might be misleading to use the calculated half-lives in river water to provide explanation for seawater. Therefore, for clarity sake, we focused only on river water study.

4. Conclusions

This study has reported the photodegradation of two carbamate insecticides, carbaryl and methomyl, in river water under environmentally-relevant conditions. To the best of our search, this is the first report to do so. Indirect photolysis mediated by $\cdot\text{OH}$ was primarily responsible for their degradation. Photodegradation studies in the presence and absence of scavengers of $\cdot\text{OH}$ and $^1\text{O}_2$ showed that $\cdot\text{OH}$ accounted for 63% and 62% degradation of methomyl and carbaryl, respectively. The involvement of $^1\text{O}_2$ was generally lower (about 30%) in the degradation of both compounds. Neither compound showed significant degradation under direct photolysis in ultrapure water. The reaction rate constants for the reactions of carbaryl and methomyl with $\cdot\text{OH}$, determined in this study, were approximately five orders higher than those of their reactions with $^1\text{O}_2$. Such preferentially high reaction rate constants with $\cdot\text{OH}$ supports the role of $\cdot\text{OH}$ as the primary ROS mediating the degradation of these compounds in river water. Furthermore,

Table 4

Calculated photochemical half-lives of the tested insecticides as mediated by $\cdot\text{OH}$ and $^1\text{O}_2$ in river water.

Insecticides	Half-lifetimes (d^{-1})	
	$\cdot\text{OH}$	$^1\text{O}_2$
Methomyl	13.30 ± 0.24^a	$>186^a$
Carbaryl	3.60 ± 0.14^b	17.0 ± 0.34^b

*Statistical comparisons were made among treatments within a single column.

*The different letters represent significant differences using Fisher's LSD test at $P \leq 0.05$.

*Each mean value came from three replicates.

kinetic estimations of degradation rate constants, calculated from reaction rate constants and ROS steady-state concentrations, agrees with experimentally-determined degradation rate constants for $\cdot\text{OH}$ -mediated degradation. This validates the experimental observation and confirms the importance of $\cdot\text{OH}$ in the degradation of the compounds in river water. The degradation half-lives of the insecticides as mediated by $\cdot\text{OH}$ were 3.6 and 13.3 days for carbaryl and methomyl, respectively.

Declaration of competing interest

The authors declare that there is no conflict of interest regarding this publication.

Acknowledgments

The Japan Society for the Promotion of Science (JSPS) is acknowledged for funding an invitation research fellowship provided to Professor Aly Derbalah at Hiroshima University, Japan. This study was supported by JSPS KAKENHI grant number 16KT0149 and 161T05622, Japan. We thank Alexander Humphreys, PhD, Jeremy Kamen, MSc, and Amanda MacMillan, PhD, from Edanz Group (www.edanzediting.com/ac) for editing a draft of this manuscript.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.chemosphere.2019.125464>.

References

- Alhousari, F., 2011. Fate and Behavior of Acidic Rice Herbicides in Lagoon Waters of Camargue (Rhône River Delta, France) Université De Provence Aix- Marseille I. PhD Thesis, p. 4. France.
- Alhousari, F., Vione, D., Chirona, S., Barbati, S., 2010. Reactive photoinduced species in estuarine waters. Characterization of hydroxyl radical, singlet oxygen and dissolved organic matter triplet state in natural oxidation processes. *Photochem. Photobiol. Sci.* 9, 78–86.
- Arakaki, T., Miyake, T., Shibata, M., Sakugawa, H., 1999. Photochemical formation and scavenging of hydroxyl radical in rain and dew waters. *Nippon Kagaku Kaishi* 5, 335–340 [in Japanese].
- Arakaki, T., Hamdun, A.M., Uehara, M., Okada, K., 2010. Photochemical formation of hydroxyl radicals in red soil-polluted seawater on the North of Okinawa Island, Japan. *Wat. Air and Soil Pollut* 209, 191–198.
- Armbrust, K.L., 2000. Pesticide hydroxyl radical rate constants: measurements and estimates of their importance in aquatic environments. *Environ. Toxicol. Chem.* 19 (9), 2175–2180.
- Barceló, D., Chiron, S., Fernández-Alba, A.R., Valverde, A., Alpendurada, M.F., 1996. In: Meyer, M.T., Thurman, E.M. (Eds.), *Monitoring Pesticides and Metabolites in Surface Water and Ground Water in Spain*. American Chemical Society, p. 237.
- Bazrafshan, A.A., Ghaedi, M., Rafiee, Z., Hajati, S., Ostovan, A., 2017. Nano-sized molecularly imprinted polymer for selective ultrasound-assisted micro-extraction of pesticide carbaryl from water samples: spectrophotometric determination. *J. Coll. and inter. Sci.* 498, 313–322.
- Bekdeser, B., Ozyurek, M., Guclu, K., Apak, R., 2012. Novel spectroscopic sensor for the hydroxyl radical scavenging activity measurement of biological samples. *Talanta* 99, 689–696.
- Buxton, G.V., Greenstock, C.L., Helman, W.P., Ross, A.B., 1988. Critical review of rate constants for reactions of hydrated electrons, hydrogen atoms and hydroxyl radicals ($\cdot\text{OH}$, O^-) in aqueous solutions. *J. Phys. Chem. Ref. Data* 17, 513–886.
- Charbouillot, T., Brigante, M., Mailhot, G., Maddigapu, P.R., Minero, C., Vione, D., 2011. Performance and selectivity of the terephthalic acid probe for $\cdot\text{OH}$ as a function of temperature, pH and composition of atmospherically relevant aqueous media. *J. Photochem. Photobiol., A* 222, 70–76.
- Chattoraj, S., Mondal, N.K., Das, B., Roy, P., Sadhukhan, B., 2014. Carbaryl removal from aqueous solution by Lemna major biomass using response surface methodology and artificial neural network. *J. of Environ. Chem. Engin.* 2, 1920–1928.
- Chowdhury, M.A., Banik, S., Uddin, B., Moniruzzaman, M., Karim, N., Gan, S.H., 2012. Organophosphorus and carbamate pesticide residues detected in water samples collected from paddy and vegetable fields of the Savar and Dhamrai Upazilas in Bangladesh. *Int. J. Environ. Res. Public Health* 33, 18–29.
- Clark, D.J., 1994. *Physical-Chemical Properties and Environmental Fate of Pesticides*. Environmental Hazards Assessment Program, State of California. Environmental Protection Agency, California, p. 56.

- Dell'Arciprete, M.L., Santos-Juanes, L., Arques, A., Vercher, R.F., Amat, A.M., Furlong, J.P., Mártire, D.O., Gonzalez, M.C., 2010. Reactivity of neonicotinoid pesticides with singlet oxygen. *Catal. Today* 151, 137–142.
- Derbalah, A.S., Ismail, A.A., Shaheen, S.M., 2013. Monitoring of organophosphorus pesticides and remediation technologies of the frequently detected compound (chlorpyrifos) in drinking water. *Pol. J. Chem. Technol.* 15 (3), 25–34.
- Derbalah, A.S., Ismail, A.A., Hamza, A.M., Shaheen, S.M., 2014. Monitoring and remediation of organochlorine residues in water. *Water Environ. Res.* 86 (7), 584–593.
- Derbalah, A.S., Ismail, A.A., Shaheen, S.M., 2016. The presence of organophosphorus pesticides in wastewater and its remediation technologies. *Environ. Eng. & Manag. J.* 15 (8), 1777–1787.
- Diamond, M.L., Priemer, D.A., Law, N.L., 2001. Developing a multimedia model of chemical dynamics in an urban area. *Chemosphere* 44, 1655–1667.
- Draper, W.M., Crosby, D.G., 1984. Solar photooxidation of pesticides in dilute hydrogen peroxide. *J. Agric. Food Chem.* 32, 231–237.
- Drea, A.A., Naman, S.N., Jaffer, B.R., 2012. Theoretical degradation study of methomyl. *J. Appl. Chem.* 1, 126–137.
- EPA 2004. Reregistration Eligibility Decision for Carbaryl US Environmental Protection Agency, (Washington, DC).
- Grandbois, M., Latch, D.E., McNeil, K., 2008. Microheterogeneous concentrations of singlet oxygen in natural organic matter isolate solutions. *Environ. Sci. Technol.* 24, 9184–9190.
- Haag, R.W., Hoigné, J., 1986. Singlet oxygen in surface waters. 3. Photochemical formation and steady-state concentrations in various types of waters. *Environ. Sci. Technol.* 20, 341–348.
- Haag, W.R., Hoigne, J.R., Gassman, E., Braun, A.M., 1984. Singlet oxygen in surface waters — Part I: furfuryl alcohol as a trapping agent. *Chemosphere* 13 (5), 631–640.
- Kaonga, C.C., Takeda, K., Sakugawa, H., 2016. Concentration and degradation of alternative biocides and an insecticide in surface waters and their major sinks in a semi-enclosed sea, Japan. *Chemosphere* 145, 256–264.
- Katagi, T., 2018. Direct photolysis mechanism of pesticides in water. *J. Pestic. Sci.* 43, 57–72.
- Larson, S.J., Capel, D.P., Majewski, S.M., 1997. Pesticides in surface waters: distribution, trends, and governing factors. In: Gilliom, R.J. (Ed.), *Series of Pesticides in Hydrologic System*, vol. 3. Ann Arbor Press, Inc., Chelsea, Michigan, p. 373.
- Latch, D.E., McNeil, K., 2006. Microheterogeneity of singlet oxygen distributions in irradiated humic acid solutions. *Science* 311, 1743–1747.
- Lian, L., Yao, B., Hou, S., Fang, J., Yan, S., Song, W., 2017. Kinetic study of hydroxyl and sulfate radical-mediated oxidation of pharmaceuticals in wastewater effluents. *Environ. Sci. Technol.* 51 (5), 2954–2962.
- Louit, G., Hanedanian, M., Taran, F., Coffigny, H., Renault, J.P., Pin, S., 2009. Determination of hydroxyl rate constants by a high-throughput fluorimetric assay: towards a unified reactivity scale for antioxidants. *Analyst* 134, 250–255.
- Mabury, S.A., Crosby, D.G., 1996. Pesticide reactivity toward hydroxyl and its relationship to field persistence. *J. Agric. Food Chem.* 44, 1920–1924.
- McDuffie, H.H., 2001. Non-Hodgkin's lymphoma and specific pesticide exposures in men: cross Canada study of pesticides and health. *Cancer Epidemiol. Biomark. Prev.* 10, 1155–1163.
- Miller, P.L., Chin, Y.-P., 2002. Photoinduced degradation of carbaryl in a wetland surface water. *J. Agric. Food Chem.* 50 (23), 6758–6765.
- Miskoski, S., García, N.A., 1993. Influence of the peptide bond on the singlet molecular oxygen-mediated ($O_2(1\Delta g)$) photooxidation of histidine and methionine dipeptides. A kinetic study. *J. Photochem. Photobiol. B Biol.* 57, 447–452.
- Motohashi, N., Saito, Y., 1993. Competitive measurement of rate constants for hydroxyl radical reactions using radiolytic hydroxylation of benzoate. *Chem. Pharm. Bull.* 41, 1842–1845.
- Nakatani, N., 2004. Doctoral thesis, Hiroshima University (in Japanese).
- Nakatani, N., Hashimoto, N., Shindo, H., Yamamoto, M., Kikkawa, M., Sakugawa, H., 2007. Determination of photoformation rates and scavenging rate constants of hydroxyl radicals in natural waters using an automatic light irradiation and injection system. *Anal. Chim. Acta* 581 (2), 260–267.
- Peterson, B.M., McNally, A.M., Cory, R.M., Thoemke, J.D., Cotner, J.B., McNeill, K., 2012. Spatial and temporal distribution of singlet oxygen in Lake Superior. *Environ. Sci. Technol.* 46 (13), 7222–7229.
- Richard, C., Canonica, S., 2005. Aquatic phototransformation of organic contaminants induced by coloured dissolved natural organic matter. In: Boule, P., Bahnemann, D.W., Robertson, P.K.J. (Eds.), *Environmental Photochemistry Part II. The Handbook of Environmental Chemistry*, vol. 2M. Springer, Berlin, Heidelberg, pp. 299–323.
- Rodgers and Snowden, 1982. Lifetime of Oxygen ($O_2(1\Delta g)$) in liquid water as determined by time-resolved infrared luminescence measurements. *J. Am. Chem. Soc.* 104 (20), 5541–5543.
- Ruggeri, G., Ghigo, G., Maurino, V., Minero, C., Vione, D., 2013. Photochemical transformation of ibuprofen into harmful 4-isobutylacetophenone: pathways, kinetics, and significance for surface waters. *Wat. Res.* 47 (16), 6109–6121.
- Scully, F., Hoigné, J., 1987. Rate constants for reactions of singlet oxygen with phenols and other compounds in water. *Chemosphere* 16 (4), 681–694.
- Street, J.C., 1981. Pesticides and the immune system. In: Sharma, R.P. (Ed.), *Immunologic Considerations in Toxicology*. 1. CRC Press, Boca Raton FL, pp. 45–66.
- Struger, J., Grabuski, J., Cagampan, S., Sverko, E., Marvin, C., 2016. Occurrence and distribution of carbamate pesticides and metalaxyl in southern ontario surface waters. *Bull. Environ. Contam. Toxicol.* 96, 423–431.
- Takeda, K., Takedoi, H., Yamaji, S., Ohta, K., Sakugawa, H., 2004. Determination of hydroxyl radical photoproduction rates in natural waters. *Anal. Sci.* 20, 153–158.
- Takeda, K., Katoh, S., Mitsui, Y., Nakano, S., Nakatani, N., Sakugawa, H., 2014. Spatial distributions of and diurnal variations in low molecular weight carbonyl compounds in coastal seawater, and the controlling factors. *Sci. Total Environ.* 493, 454–462.
- Takeda, K., Fujisawa, K., Nojima, H., Kato, R., Ueki, R., Sakugawa, H., 2017 a. Hydroxyl radical generation with a high power ultraviolet light emitting diode (UV-LED) and application for determination of hydroxyl radical reaction rate constants. *J. Photochem. Photobiol. A Chem.* 340, 8–14.
- Takeda, K., Nojima, H., Kuwahara, K., Chidya, R.C., Adesina, A.O., Sakugawa, H., 2017 b. Nanomolar determination of hydrogen peroxide in coastal seawater based on the Fenton reaction with terephthalate. *Anal. Sci.* 34, 459–464.
- Tomlin, C., 2003. *The Pesticide Manual*, thirteenth ed. BCPC, Hampshire, UK.
- Trapp, S., Matthies, M., 1995. Generic one-compartment model for uptake of organic chemicals by foliar vegetation. *Environ. Sci. Technol.* 29, 2333–2338.
- Vermilyea, A.W., Voelker, B.M., 2009. Photo-fenton reaction at near neutral pH. *Environ. Sci. Technol.* 43, 6927–6933.
- Vione, D., Maurino, V., Minero, C., Carlotti, M.E., Chiron, S., Barbati, S., 2009. Modeling the occurrence and reactivity of the carbonate radical in surface freshwater. *C. R. Chimie* 12, 865–871.
- Vione, D., Maddigapu, P.R., De Laurentis, E., Minella, M., Pazzi, M., Maurino, V., Minero, C., Kouras, S., Richard, C., 2011. Modelling the photochemical fate of ibuprofen in surface waters. *Wat. Res.* 45, 6725–6736.
- Walse, S.S., Morgan, S.L., Kong, L., Ferry, J.L., 2004. Role of dissolved organic matter, nitrate, and bicarbonate in the photolysis of aqueous fipronil. *Environ. Sci. Technol.* 38, 3908–3915.
- Wilson, P.C., Foos, J.F., 2006. Survey of carbamate and organophosphorous pesticide export from a south Florida (U.S.A.) agricultural watershed: implications of sampling frequency on ecological risk estimation. *Environ. Toxicol. Chem.* 25, 2847–2852.
- Yang, X.F., Guo, X.Q., 2001. Fe(II)-EDTA chelate-induced aromatic hydroxylation of terephthalate as a new method for the evaluation of hydroxyl radical-scavenging ability. *Analyst* 126, 928–932.
- Zepp, R.G., Cline, M.D., 1977. Rate of direct photolysis in aquatic environment. *Environ. Sci. Technol.* 11, 359–366.
- Zepp, R.G., Baughman, G.L., Scholtzauer, P.F., 1981. Comparison of photochemical behavior of various humic substances in water: sunlight induced reactions of aquatic pollutants photosensitized by humic substance. *Chemosphere* 10, 109–177.
- Zepp, R.G., Scholtzauer, P.F., Sink, R.M., 1985. Photosensitized transformations involving energy transfer in natural waters: role of humic substances. *Environ. Sci. Technol.* 19, 74–81.