

## Editorial

# Do PCBs Modify the Thyroid-Adipokine Axis during Development?

Ahmed RG\*

Department of Zoology, Beni-Suef University, Egypt

\*Corresponding author: Ahmed RG, Department of Zoology, Division of Anatomy and Embryology, Faculty of Science, Beni-Suef University, Beni-Suef, Egypt, Tel: 002-010-9147-1828; Email: r\_g\_a\_ahmed@yahoo.com & ahmedragab08@gmail.com

Received: October 02, 2014; Accepted: October 06, 2014; Published: October 07, 2014

## Editorial

Polychlorinated Biphenyls (PCBs) are widespread global contaminants and they are a class of industrial compounds (anthropogenic chemicals) consisting of paired phenyl rings with various degrees of chlorination [1-4]. In the United States, PCBs are manufactured as complex mixtures under the trade name Aroclor with a designation indicating the extent of chlorination [5]. Also, the U.S. Environmental Protection Agency recognizes PCBs as unavoidable toxins (2010 21CFR109.30). They are highly lipophilic organochlorines, accumulated in the food chain and bioaccumulated in the biological tissues (adipose tissue) to levels that are associated with a variety of health effects in human and animals [6]. These potential effects are amplified during the development [7] and their elimination half lives are about 6 to 10 years [8].

PCBs may affect the growth and development by impairing thyroid function [9]. Interestingly, their interference mechanism with Thyroid Hormones (THs) metabolism in the mother, fetus and newborn may injure the developing brain [10], even at background environmental levels [11], via changing the availability of Thyroid Receptors (TRs) [12], deiodinases [13] and TH transporters [14], or altering GABA-mediated pathways [15], monoaminergic pathways and the activity of cholinesterases [10]. Also, these changes may mediate by Aryl Hydrocarbon Receptors (AhRs) [16]. In my recent work [17], I observed that the administration of 2,3,6-2',5'-Pentachlorinated Biphenyl (PCB 95) induced a reduction ( $P<0.01$ ) in sera Thyroxine (T4), Triiodothyronine (T3) and Growth Hormone (GH) levels, and an increase ( $P<0.01$ ) in serum Thyroid Stimulating Hormone (TSH) level at Postnatal Days (PNDs) 17 and 18. At PND 18, the administration of PCB 95 led to thyroid dysgenesis and the thyroid gland exhibited some histopathological changes, such as follicular destruction, luminal obliteration, edema, interfollicular fibroblast proliferation, hemorrhage, and hypertrophy with reduced colloidal contents. This dysmorphogenesis and dysgenesis may be attributed to the increase in the level of DNA-fragmentation at PNDs 17 and 18. These results imply that PCB 95 may act as disruptor for the developmental Hypothalamic-Pituitary-Thyroid Axis (HPTA). Obviously, exposure to PCB induced hypothyroidism in rats [18], in sledge dogs [19], in monkeys [20] and in children living near PCB-contaminated sites [21]. Also, a PCBs [7] and PCB metabolites

[22] may act directly on TR to modulate its action or indirectly on an unknown TR-binding protein, which may then conformational changes in TR-DNA binding domain to dissociate TR from T3-Responsive Element (TRE). Indeed, the Persistent Organic Pollutants (POPs) show resistance to biological degradation and may exert a large variety of adverse health effects [23].

On the other hand, in animal and human studies, PCBs appear to disrupt the levels of adiponectin [23], leptin [24], tumor necrosis factor (TNF- $\alpha$ ) [25], insulin [26], body weight and fat depots [27]. In my recent work [17], the hypothyroid state due to PCB 95 revealed a higher ( $P<0.01$ ) levels of sera leptin, adiponectin and TNF- $\alpha$ , and a lower ( $P<0.01$ ) levels of sera Insulin Growth Factor (IGF-I) and insulin in both PNDs with respect to control group. In particular, the administration of PCB 95 at PND 18 led to a maximal increase in the concentration of adiponectin and the highest drop in the concentrations of insulin and IGF1, respectively. This is probably due to the disturbance of the hormonal homeostatic mechanisms on this day. Interestingly, the body weight of the neonates in the PCB 95 group exhibited severe decreases throughout the experimental period in relation to that of the control group. The hypothyroidism by PCB 95 could impair the adipokines axis, fat metabolism and in general the postnatal development. Similarly, the full mixture of PCB, organochlorine pesticides and methylmercury also moderately increased (in an additive fashion) hypothalamic levels of the TNF- $\alpha$  and interleukins (IL6 or IL12) [28]. Also, Baker et al. [29] undertook that the dysfunction in the glucose homeostasis by PCBs could be AhR mediated. A previous laboratory experiments showing that a lower dose of Aroclor 1254 (200  $\mu\text{g}/\text{kg}/\text{day}$ ) [27] or a higher dose of bisphenol A (BPA) [30] induced a significantly decreased rat body weight, fat depots and a lower feeding efficiency. Generally, PCBs produce a wide spectrum of toxic effects in animals including body weight loss, toxicity, and teratogenicity [31]. However, a negative correlation was found between the level of adiponectin and PCB 153 [23], PCB 77 (3,3',4,4'-tetrachlorobiphenyl) [25], 2,3,7,8-Tetrachlorodibenzo-P-Dioxin (TCDD) [32] and BPA [33]. More importantly, the changes in the level of serum adiponectin play a modest role in thyroid dysfunction in human [34].

From this point of view, the endocrine disrupting compounds can produce complex, mosaic effects during the life cycle [35]. Conclusively, loss of body weight may suggest a decline in the general health level of animals [36], which can be important in the interpretation of thyroid effects. However, these mechanisms associated with developing thyroid dysfunction are not fully understood.

## References

1. Murk AJ, Rijntjes E, Blaauboer BJ, Clewell R, Crofton KM, Dingemans MM, et al. Mechanism-based testing strategy using *in vitro* approaches for identification of thyroid hormone disrupting chemicals. *Toxicol In Vitro*. 2013; 27: 1320-1346.

2. Arukwe A, Olufsen M, Cicero N, Hansen MD. Effects on development, growth responses and thyroid-hormone systems in eyed-eggs and yolk-sac larvae of Atlantic salmon (*Salmo salar*) continuously exposed to 3,3',4,4'-tetrachlorobiphenyl (PCB-77). *J Toxicol Environ Health A*. 2014; 77: 574-586.
3. Quinete N, Schettgen T, Bertram J, Kraus T. Analytical approaches for the determination of PCB metabolites in blood: a review. *Anal Bioanal Chem*. 2014; 406: 6151-6164.
4. Naveau E, Pinson A, Gérard A, Nguyen L, Charlier C, Thomé JP, et al. Alteration of rat fetal cerebral cortex development after prenatal exposure to polychlorinated biphenyls. *PLoS One*. 2014; 9: e91903.
5. Karcher SC, Small MJ, VanBriesen JM. Statistical method to evaluate the occurrence of PCB transformations in river sediments with application to Hudson River data. *Environ Sci Technol*. 2004; 38: 6760-6766.
6. Miller VM, Sanchez-Morrissey S, Brosch KO, Seegal RF. Developmental coexposure to polychlorinated biphenyls and polybrominated diphenyl ethers has additive effects on circulating thyroxine levels in rats. *Toxicol Sci*. 2012; 127: 76-83.
7. Koibuchi N, Iwasaki T. Regulation of brain development by thyroid hormone and its modulation by environmental chemicals. *Endocr J*. 2006; 53: 295-303.
8. Norström K, Czub G, McLachlan MS, Hu D, Thorne PS, Hornbuckle KC. External exposure and bioaccumulation of PCBs in humans living in a contaminated urban environment. *Environ Int*. 2010; 36: 855-861.
9. Kobayashi K, Miyagawa M, Wang RS, Suda M, Sekiguchi S, Honma T. Effects of in utero exposure to 2,2',4,4',5,5'-hexachlorobiphenyl on postnatal development and thyroid function in rat offspring *Industrial Health*. 2009; 47: 189-197.
10. Ahmed RG. Perinatal TCDD exposure alters developmental neuroendocrine system. *Food Chem Toxicol*. 2011; 49: 1276-1284.
11. Darras VM. Endocrine disrupting polyhalogenated organic pollutants interfere with thyroid hormone signalling in the developing brain. *Cerebellum*. 2008; 7: 26-37.
12. Zoeller RT, Crofton KM. Thyroid hormone action in fetal brain development and potential for disruption by environmental chemicals. *Neurotoxicology*. 2000; 21: 935-945.
13. Morse DC, Groen D, Veerman M, van Amerongen CJ, Koëter HB, Smits van Prooije AE, et al. Interference of polychlorinated biphenyls in hepatic and brain thyroid hormone metabolism in fetal and neonatal rats. *Toxicol Appl Pharmacol*. 1993; 122: 27-33.
14. Friesema EC, Visser WE, Visser TJ. Genetics and phenomics of thyroid hormone transport by MCT8. *Mol Cell Endocrinol*. 2010; 322: 107-113.
15. Bushnell PJ, Rice DC. Behavioral assessments of learning and attention in rats exposed perinatally to 3,3',4,4',5-pentachlorobiphenyl (PCB 126) *Neurotoxicol Teratol*. 1999; 21: 381-392.
16. Fisher JW, Campbell J, Muralidhara S, Bruckner JV, Ferguson D, Mumtaz M, et al. Effect of PCB 126 on hepatic metabolism of thyroxine and perturbations in the hypothalamic-pituitary-thyroid axis in the rat. *Toxicol Sci*. 2006; 90: 87-95.
17. Ahmed RG. Early weaning PCB 95 exposure alters the neonatal endocrine system: thyroid adipokine dysfunction. *J Endocrinol*. 2013; 219: 205-215.
18. Powers BE, Widholm JJ, Lasky RE, Schantz SL. Auditory deficits in rats exposed to an environmental PCB mixture during development. *Toxicol Sci*. 2006; 89: 415-422.
19. Kirkegaard M, Sonne C, Dietz R, Letcher RJ, Jensen AL, Stige HS, et al. Alterations in thyroid hormone status in Greenland sledge dogs exposed to whale blubber contaminated with organohalogen compounds. *Ecotoxicol Environ Saf*. 2011; 74: 157-163.
20. Boas M, Feldt-Rasmussen U, Main KM. Thyroid effects of endocrine disrupting chemicals. *Mol Cell Endocrinol*. 2012; 355: 240-248.
21. Schell LM, Gallo MV, Decaprio AP, Hubicki L, Denham M, Ravenscroft J, Akwesasne Task Force on the Environment. Thyroid function in relation to burden of PCBs, p,p'-DDE, HCB, mirex and lead among Akwesasne Mohawk youth: a preliminary study. *Environ Toxicol Pharmacol*. 2004; 18: 91-99.
22. Miyazaki W, Iwasaki T, Takeshita A, Tohyama C, Koibuchi N. Identification of the functional domain of thyroid hormone receptor responsible for polychlorinated biphenyl-mediated suppression of its action *in vitro*. *Environ Health Perspect*. 2008; 116: 1231-1236.
23. Mullerova D, Kopecky J, Matejkova D, Muller L, Rosmus J, Racek J, et al. Negative association between plasma levels of adiponectin and polychlorinated biphenyl 153 in obese women under non-energy-restrictive regime. *Int J Obes (Lond)*. 2008; 32: 1875-1878.
24. Nieminen P, Hyvärinen H, Käkälä R, Asikainen J. Plasma leptin and thyroxine of mink (*Mustela vison*) vary with gender, diet and subchronic exposure to PCBs. *Comp Biochem Physiol A Mol Integr Physiol*. 2000; 127: 515-522.
25. Arsenescu V, Arsenescu RI, King V, Swanson H, Cassis LA. Polychlorinated biphenyl-77 induces adipocyte differentiation and proinflammatory adipokines and promotes obesity and atherosclerosis. *Environ Health Perspect*. 2008; 116: 761-768.
26. Everett CJ, Frithsen I, Player M. Relationship of polychlorinated biphenyls with type 2 diabetes and hypertension. *J Environ Monit*. 2011; 13: 241-251.
27. Anbalagan J, Kanagaraj P, Srinivasan N, Aruldas MM, Arunakaran J. Effect of polychlorinated biphenyl, Aroclor 1254 on rat epididymis. *Indian J Med Res*. 2003; 118: 236-242.
28. Hayley S, Mangano E, Crowe G, Li N, Bowers WJ. An *in vivo* animal study assessing long-term changes in hypothalamic cytokines following perinatal exposure to a chemical mixture based on Arctic maternal body burden *Environ Health*. 2011; 10: 65.
29. Baker NA, Karounos M, English V, Fang J, Wei Y, Stromberg A, et al. Coplanar polychlorinated biphenyls impair glucose homeostasis in lean C57BL/6 mice and mitigate beneficial effects of weight loss on glucose homeostasis in obese mice. *Environ Health Perspect*. 2013; 121: 105-110.
30. Nunez AA, Kannan K, Giesy JP, Fang J, Clemens LG. Effects of bisphenol A on energy balance and accumulation in brown adipose tissue in rats. *Chemosphere*. 2001; 42: 917-922.
31. Safe S. Polychlorinated biphenyls (PCBs) and polybrominated biphenyls (PBBs): biochemistry, toxicology, and mechanism of action. *Crit Rev Toxicol*. 1984; 13: 319-395.
32. Kern PA, Said S, Jackson WG Jr, Michalek JE. Insulin sensitivity following agent orange exposure in Vietnam veterans with high blood levels of 2,3,7,8-tetrachlorodibenzo-p-dioxin. *J Clin Endocrinol Metab*. 2004; 89: 4665-4672.
33. Kidani T, Kamei S, Miyawaki J, Aizawa J, Sakayama K, Masuno H. Bisphenol A downregulates Akt signaling and inhibits adiponectin production and secretion in 3T3-L1 adipocytes. *J Atheroscler Thromb*. 2010; 17: 834-843.
34. Díez JJ, Iglesias P. The role of the novel adipocyte-derived hormone adiponectin in human disease. *Eur J Endocrinol*. 2003; 148: 293-300.
35. Zoeller RT, Brown TR, Doan LL, Gore AC, Skakkebaek NE, Soto AM, et al. Endocrine-disrupting chemicals and public health protection: a statement of principles from The Endocrine Society. *Endocrinology*. 2012; 153: 4097-4110.
36. Fernandes GS, Arena AC, Fernandez CD, Mercadante A, Barbisan LF, Kempinas WG. Reproductive effects in male rats exposed to diuron. *Reprod Toxicol*. 2007; 23: 106-112.