

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/319528365>

LABORATORY BIOASSAY OF METARHIZTUM ANISO₂LIAE (METSCH.) AGAINST THE TWO SPOTTED SPIDER MITE/ TETRANYCHUS URTICAE KOCH (ACARI: TETRANYCHIDAE)

Conference Paper · November 2005

CITATIONS

0

READS

15

1 author:



Rania AHMED Abd El-Wahab

Agricultural Research Center, Egypt

36 PUBLICATIONS 51 CITATIONS

SEE PROFILE

Some of the authors of this publication are also working on these related projects:



Pesticides' Resistance Solution [View project](#)



Pesticides [View project](#)




**Arab Republic of Egypt
Ministry of Agriculture
and Land Reclamation**

EGYPTIAN JOURNAL OF AGRICULTURAL RESEARCH

**Special Issue
The Third International Conference
Of Plant Protection Research Institute
26 - 29 November 2005**

VOLUME 83, NUMBER 3

2005



LABORATORY BIOASSAY OF *METARHIZIUM ANISOPLIAE* (METSCH.) AGAINST THE TWO SPOTTED SPIDER MITE, *TETRANYCHUS URTICAE* KOCH (ACARI: TETRANYCHIDAE)

RANIA A. ABD EL WAHAB

Plant protection Research institute, Dokki, Giza, Egypt

(Manuscript received 4 October 2005)

Abstract

Laboratory bioassay of an entomopathogenic fungi, *Metarhizium anisopliae* (Metsch.) Sorokin, which maintained on potato dextrose agar (PDA) Difco, and liquid medium was evaluated against the two spotted spider mite, *Tetranychus urticae* Kock adult females. This bioassay revealed that liquid medium supernatants (LMS) caused mortality up to 90% while *M. anisopliae* spores resulted, from PDA (solid medium) caused 76.67% mortality at 8×10^5 spores/ml.

EC₅₀ and EC₉₀ values of *M. anisopliae* spores were 3.38×10^5 and 16.62×10^5 spores/ml, respectively.

Key words: Bioassay, *Metarhizium anisopliae*, *Tetranychus urticae*

INTRODUCTION

Available knowledge of mite diseases is increasing in recent years because of the economic importance of many mite species especially eriophyoids and spider mites. Fungal pathogens often an important role in the regulation of natural mite populations (Van Der Geest et al., 2000).

Tetranychid mite, *Tetranychus urticae* can be infested by entomophthorales fungi, such as Neozygites, which possess a restricted host spectrum and are obligatory pathogens. Also, deuteromycetes are able to kill *T. urticae* even by true spores or mycotoxins which often cause host death prior to complete colonization of the body (Hajek, 1997).

Metarhizium is a clear example of Deuteromycetes fungi. The first species of the genus metarhizium (Subdivision. Deuteromycotina; Class Hyphomycetes; Order Moniliales), *Metarhizium anisopliae*, was isolated from the coleopteran species, *Anisopliae austriacae* by Metachinkoff in 1878. *Metarhizium anisopliae* can produce three spore types in vitro. Aerial conidia in solid culture and both of blastospores and submerged conidia in liquid culture (Jenkins and Prior, 1993). This fungi improved its efficacy against many insects such as grasshoppers and locusts (Lomer et al., 2001)

and mites. *M. anisopliae* provided an excellent control to varroa mites infested honeybees (Flores,2004). Most infested mites died within 3-5 days without any bad hazards to bees.

M. anisopliae hyphae could easily penetrate the soft unsclerotised exoskeleton of astigmatid sheep scab mite, *Psorptes ovis* and caused 77% mortality of mites which were exposed to 1×10^8 conidia/ml at 25°C (Brooks et al., 2003).

Moreover, *M. anisopliae* produce many toxins, the most important group named destruxins which can be used as pesticides (Glover, 2005). Destruxins has a lack of mammalian toxicity at high exposure levels. Beside that the quantities of toxic secondary metabolites produced by fungi in vivo were much less than those in nutrient rich liquid media. Therefore, use of mycoinsecticides was not expected to result in harmful levels to the environment (EPA, 2005). So, *M. anisopliae* is generally safe to natural enemies.

This research aimed to evaluate the lab. efficacy of *M. anisopliae* spores and *M.anisopliae* liquid medium supernatant (LMS) against *Tetranychus urticae* adult females.

MATERIALS AND METHODS

A)Fungal medium

- a) **Solid media:** *Metarhizium anisopliae* spores were isolated from naturally infested *Tetranychus urticae* adult females. Spores were maintained on PDA (Potato Dextrose Agar) supplemented with 5% yeast extract (Difco) which had been sterilized, for 20 min at 121°C. To prepare fungal inocula conidia from 2-3 weeks old cultures (20 + 2.70°C and 70%RH) were scraped from the surface of the plates with a sterile scalpe and suspended in 0.05% aqueous Tween 80. Haemocytometer was used to estimate the conidial concentration and subsequent appropriate dilutions were made.

A)Fungal medium

- b) **Solid media:** *Metarhizium anisopliae* spores were isolated from naturally infested *Tetranychus urticae* adult females. Spores were maintained on PDA (Potato Dextrose Agar) supplemented with 5% yeast extract (Difco) which had been sterilized, for 20 min at 121°C. To prepare fungal inocula conidia from 2-3 weeks old cultures (20 + 2.70°C and 70%RH) were scraped from the surface of the plates with a sterile scalpe and suspended in 0.05% aqueous Tween 80. Haemocytometer was used to estimate the conidial concentration and subsequent appropriate dilutions were made.

summer and on castor oil leaves at winter. This colony was left for one year under the previous conditions in order to get a homogenous and sensitive colony.

C) Laboratory bioassay: *Metarhizium anisopliae* was bioassayed using an adulticidal technique (Edge and James, 1982). Briefly, 20 young adult female mites were transferred to castor oil leaf discs (2cm diameter). Mites on the leaf discs were then sprayed with aqueous suspensions of *M. anisopliae* spores and with 10 ml (LMS) of each spores concentration. Controls of *M. anisopliae* spores treatment were sprayed with water + 0.05% Tween 80 but in the case of (LMS) treatments, controls sprayed with water only. Sprayed leaf discs (three per concentration) were maintained under constant light on moistened cotton wool for 72h (25±2°C, 60±5%RH). Mortality percentages were determined and corrected by using Abbott's formula (1925). Data were analysed using a probit program written in GENSTAT5 statistical software (Barchia, 2001). LC₅₀ and LC₉₀ values plus their 95% fiducial limits, were calculated using the probit method outlined in (Finney, 1971).

RESULTS AND DISCUSSION

Data showed that adult females of *T. urticae* were affected by serial concentrations of *M. anisopliae* spores and its liquid media supernatants (LMS).

Mortality percentage caused by fungi spores of mite females at high concentration 8×10^5 spores/ml was 76.67%. Table (1), but it was increased to 93.33% in the case of (LMS). Table (2).

Estimates of the EC₅₀ and EC₉₀ with 95% fiducial limits computed for *M. anisopliae* spores against *T. urticae* were 3.38×10^5 (5.61×10^5 - 2.05×10^5) spores/ml and 16.2×10^5 (27.59×10^5 - 10.0×10^5) spores/ml, respectively. Table (1).

Metarhizium anisopliae as an entomopathogenic fungus improved its pathogenicity against *T. urticae* adult females in this research as shown previously.

The basic infective unit of fungus was spore which had to adhere to the cuticle of host and germinate. Attachment, the first step of infection process, was mediated by chemical components in the outer layers of the spore and the metal cuticle (Parker et al., 2000). *M. anisopliae* hydrophobic spores were covered by proteins, lipoproteins and polysaccharides which helped spores to adhere themselves to the host beside that fungus secreted N-acetylglucosaminidase which concentrated on the spores surfaces when they touched hosts body, immediately (St Leger et al., 1991).

After that, spores germination took place, fastly within few hours. In the case of *T.urticae* adult females fungi infected them at moisturized ares (Charnley et al., 1997). Such as mouth parts, secretion pores (excretory proe, etc), and natural pores in tetranychid mite's such as the couple spiracles which located on ganthosoma, joints of legs, chelicerae, and pedipalps and where the cuticle is less sclerotised plus humidated conditions.

The next step of infection process was penetration. *M.anisopliae* produced appressoria to help it to stuck in vitro on hard hydrophobic surfaces like cuticle. (St Leger et al., 1989) and cuticle degrading enzymes such as lipases, proteinases, and chitinases.

Once fungi reached hemoceol, it encountered the host's immune system, by secreted toxins. Then host died and it was good resource of infection of fungi.

(Nugroho and Ibrahim, 2004) investigated the effect of *M.anisopliae* against broad mite (Polyphagotarsenomus *latus* Bank) and they found that EC_{50} was 2.77×10^7 conidia/ml.

Beside that several scientists prefered using entomophthorales fungi as control agents of *T.urtica* (Leit et al., 2000) more than deuteromycetes fungi (Tamai et al., 1998). On the other hand, data concerning this research not agree with them generally, but if there was a comparison between the ability of pathogenicity of *M.anisopliae* spores and its (LMS), the last one could be the superior against *T.urticae*.

Concerning the highly good response of treatment with *M.anisopliae* (LMS), it could be explained upon two points.

The most important one depend on mycotoxins which secreted by fungi as a secondary metabolite products caused mite death prior to complete colonization of the body (Van Der Geest et al., 2000). These included cyclic depsipeptides; Cytochalsins and Destruxins (DTXs) (Clarkson and Charnely, 1996). DTXs showed highly effect as an insecticidal when ingested by mouth.

DTXs had diverse modes of action including depolarization of muscle membrane Ca^{+2} channels, inhibition hemocyte (Blood cells) function and inhibition of vascular – type ATP ase. This can be an answer of the main question: Why (LMS) of *M.anisopliae* had a fast reponse against *T.urticae* comparable with treatments by *M. anisopliae* spores only.

Secondly, blastospores which were only produced in liquid medium, thought to have faster germination rates than conidia. So it contributed in fast infection and

mortality. (Thomas et al., 1987). Spores with faster germination rates might have a greater potential for infection by reducing potential for desiccation, effects of other organisms, or less during molting while on the insect cuticle, (Dillon and Charnely 1985). Beside blastospores in liquid media there were submerged conidia, which were hydrophilic and had similar characteristics of aerial conidia (Jenkins and Prior 1993).

REFERENCES

1. Abbott, W.S. 1925. A method for computing the effectiveness of an insecticide. J. Econ. Entomol. 18: 265-267.
2. Barchia, I. 2001. Probit analysis and fiducial limits in Genstat. Mecure Resort, Gold Cost, Australia, P. 30 31stJan.-2ndFeb., 2001.
3. Brooks, A.J., M. Aquino de Muro, D. Moore and R. Wall. 2003. The control of Psoroptes mites with a fungal pathogen. Irish Soc. Parasitology and British Association of veterinary parasitologists joint Meeting. York, England. 2nd -3rd Sept. 2003.
4. Charnely, A.K., B.Cobb and J.M. Clarkson, 1997. Towards the improvement of fungal insecticides. PP:115-126. In: Microbial insecticides: Novelty or Necessity?. H.F. Evans (ed.) proc. BCPC Symposium Coventry, U.K. 16th-18th April. 1997.
5. Clarkson, J.M. and A.K. Charnely. 1996. New insights into the mechanism of fungal Pathogenesis in insects. Trends in Microbiol. 4(5): 199-206.
6. Dillon, R.J. and A.K. Charnley, 1985. A technique for accelerating and synchronising germination of conidia of the entomopathogenic fungus, *Metarhizium anisopliae*. Arch. Micro biol, 142: 204-206.
7. Edge, V.E. and D.G. James. 1982. Detection of Cyhexatin resistance in two spotted mite, *Tetranychus urtica* koch (Acari: Tetranychidae). J.Austral. Entomol. Soc. 21:198.
8. EPA (U.S.Environmental Protection Agency). 2005. Biopesticides registration action documents, *Metarhizium anisopliae* newly reports.
9. Finney, D.J. 1971. Probit Analysis (Third Edition). Cambridge Univ. Press, Cambridge
10. Flores, A. 2004. Saving bees: Fungus Found to attack varroa mites. The Buzz New letter, Nov. 14.

11. Glober, J.B. 2005. Handbook of secondary fungal metabolites. J. Nat. Prod. 68(1): 150-151.
12. Hajek, A.E. 1997. Ecology of terrestrial fungal entomopathogens. Adv. Microbial Ecol. 15: 193-249.
13. Jenkins, N. E. and C. Prior 1993. Growth and formation of true conidia by *Metarhizium flavoviridae* in simple liquid medium . Mycol. Res. 97: 1489-1494.
14. Leite, L.G, L. Smith and D.W, Robert. 2000. In Vitro production of hyphal bodies of the mite pathogenic fungi, *Neozygites floridana*. Mycologia, 92: 201-207.
15. Lomer, C.J.,R.P. Bateman, J. Langewald and M.B. Thomas. 2001. Biological control of locusts and grasshoppers. Ann. Rev. Entomol . 46: 667-702
16. Nugroha, I. and Y. Ibrahim 2004. Laboratory Bioassay of some Entomopathogenic fungi against broad mite, *Polyphagotarsonemus latus* Bank. Int. J. Agric. Bio (2): 223-225.
17. Parker, B.L.,M. Skinner, M. Brownbridge and M. ElBoushesini. 2000. Control insect pests with entomopathogenic fungi. Symposium on biological control of insect pests. Seventh Arab Congress of protection, Amman Jordan. 22-26, Oct. 2001.
18. St Leger, R.J., D.W. Roberts and R.C. Staples 1991. A model to explain differentiation of appressoria by germlings of *Metarhizium anisopliae*. J. Invert. Pathol. 57: 299-310.
19. St Leger, R.J., T.M. Butt, M.S. Staples and D.W. Roberts 1989. Production in vitro of appressoria by an Entomopathogenic fungi, *Metarhizium anisopliae*. Exp. Mycol. 13: 274-288.
20. Tamai, M.A., B. Alves, R.B. Lopes and P.S. Neves, 1998. Avaliacao de fungo entomopatogenicos para a controle de *Tetranychus urticae* Koch. Rio de Janeiro, Brazil. P. 1066.
21. Thomas, M.B. , S.N. Wood and C.J. Lomer. 1997. Persistence of *Metarhizium flavoviridae* and consequences for biological control of grasshoppers and locusts. Pesticide Sci. 49: 47-55.
22. Van Der Geest, L.P.S, S.L. Elliot, J.A. Breeuwer and E.A. Beerling. 2000 Diseases of mites. Exp. Appl. Acarol. 24: 497-560.

Table1. Results of probit analysis of *Metarhizium anisopliae* spores against *Tetranychus urticae* Koch adult females

Conc. Spores/ml	% mortality	EC ₅₀		EC ₉₀		Slope
		Value	95% fiducial limits	Value	95% fiducial limits	
1x10 ⁵	16.67	3.38	5.61 – 2.05	16.2	27.9-10.01	1.85
2x10 ⁵	33.33					
4x10 ⁵	53.33					
8x10 ⁵	76.67					

Table 2. Effect of *Metarhizium anisopliae* liquid media supernatants against *Tetranychus urticae* koch adult females.

Spores/ml water/100ml liquid media	% Mortality
1x10 ⁵	30
2x10 ⁵	50
4x10 ⁵	66.67
8x10 ⁵	93.33